DOI: 10.1111/1365-2745.70082

RESEARCH ARTICLE

Inbreeding depression, functional traits and phenotypic plasticity in an endangered tree species with a mixed mating system

Jean Pierre Ngongo^{1,2,3} | Chadrack Kafuti^{4,5,6} | Hans Beeckman⁵ | Léopold Ndjele² | Espoir Ibrahim² | Crispin Ilunga-Mulala^{7,8} | Hulda Hatakiwe⁹ | Yves Brostaux¹⁰ | Dieu-Merci Assumani Angbonda^{3,11,12} | Thomas Drouet¹³ | Nils Bourland^{5,9,14} | Olivier J. Hardy³

¹Department of Natural and Renewable Resources Management, Faculty of Agricultural Sciences, University of Kindu, Kindu, Democratic Republic of the Congo; ²Faculty of Sciences, University of Kisangani, Kisangani, Democratic Republic of the Congo; ³Evolutionary Biology and Ecology, CP 160/12, Faculty of Sciences, Université Libre de Bruxelles, Brussels, Belgium; ⁴UGCT-Woodlab-UGent, Laboratory of Wood Technology, Ghent University, Ghent, Belgium; ⁵Service of Wood Biology, Royal Museum for Central Africa, Tervuren, Belgium; ⁶Department of Natural Resources Management, Faculty of Agricultural Sciences, University of Kinshasa, Kinshasa, Democratic Republic of the Congo; ⁷Forest Is Life, TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium; ⁸Faculté des Sciences, Université du Cinquantenaire de Lwiro, Lwiro, Democratic Republic of the Congo; ⁹Center for International Forestry Research (CIFOR), Bogor, Indonesia; ¹⁰Applied Statistics, Computer Science and Modelling Research Unit, TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium; ¹¹Faculté de Gestion des Ressources Naturelles Renouvelables, Université de Kisangani, Kisangani, Democratic Republic of Congo; ¹²Institut National Pour l'Etude et la Recherche Agronomiques de Yangambi (INERA-Yangambi), Yangambi, Democratic Republic of the Congo; ¹³Plant Ecology and Biogeochemistry, CP 244, Faculty of Sciences, Université Libre de Bruxelles, Brussels, Belgium and ¹⁴Ressources & Synergies Development, Singapore, Singapore

Correspondence

Jean Pierre Ngongo Email: fistonngongo1@gmail.com

Funding information

European Union (EU) via the Center for International Forestry Research (CIFOR); FORETS project (EEC40); Académie de Recherche et d'Enseignement Supérieur (ARES); Belgian Directorate-General for Development Cooperation and Humanitarian Aid (DGD) via PilotMAB; Belgian Fund for Scientific Research (F.R.S-FNRS)

Handling Editor: Albert Vilà-Cabrera

Abstract

- 1. Most tree species can suffer from inbreeding depression (ID), which they escape by reproducing predominantly through outcrossing. A remarkable exception is *Pericopsis elata*, an African timber species naturally producing 54% of self-fertilized seeds in the eastern Congo Basin. This species is highly logged and suffers from a deficit of natural regeneration, so that silviculture is needed for its sustainable management. While selecting good genetic material can increase the value of plantations, we lack fundamental biological knowledge on the effect of inbreeding and competition on growth potential, variability in leaf traits and phenotypic plasticity (PP). We hypothesize that ID in *P. elata* could result from the expression of deleterious mutations affecting functional traits, or from a reduction of adaptive PP in inbred genotypes.
- 2. To test our hypotheses, 540 *P. elata* seedlings were monitored for 4 years in a Nelder-type device, in which trees were planted along concentric circles to generate a density gradient. Nine leaf morphological traits (including specific leaf area, stomata density and size), eight leaf chemical traits, diameter and total height were measured regularly on 60 individuals, while paternity analyses allowed distinguishing inbred and outbred plants. To explain the observed ID on growth, we

© 2025 The Author(s). Journal of Ecology © 2025 British Ecological Society.

tested whether inbreeding affected leaf traits and/or their plasticity expressed across years, across the density gradient or across sunlight exposure.

- 3. Outbred plants grew faster than inbred ones, demonstrating ID for each level of competition. Despite the significant correlation found between specific LA and growth, and the impact of planting density, plant age and leaf exposure to sunlight on multiple traits, mean leaf trait values did not differ according to inbreeding. However, a few leaf traits (chlorophyll content, maximum stomatal water vapour conductance and leaf fresh mass) showed significantly higher plasticity in outbred than inbred plants.
- 4. Synthesis. The observed ID on growth traits was not explained by a direct effect of inbreeding on the mean values of leaf traits but possibly by a reduction in PP with inbreeding. Additional studies on the interplay between ID, functional traits and plasticity should be conducted at the intraspecific level to identify general patterns.

KEYWORDS

functional traits, inbreeding depression, mating system, Nelder device, *Pericopsis elata*, phenotypic plasticity, silviculture

1 | INTRODUCTION

Inbreeding depression (ID), the negative impact of inbreeding on fitness, is a pervasive phenomenon in plants and animals, usually attributed to the expression of deleterious recessive mutations in more homozygous genotypes (Charlesworth & Willis, 2009; Hedrick et al., 2016). The first quantification of the harmful effects of inbreeding on plants was done by Charles Darwin who showed that inbreeding reduces the vigour and fertility of plant species (Darwin, 1876). This ID can be experienced from the earliest stages of the plant's life cycle (Hufford & Hamrick, 2003) but its severity tends to rise with age (Hardner & Potts, 1995). Harmful effects of inbreeding include a reduction of the seed production (Herlihy & Eckert, 2002), germination rates (Amorim et al., 2023; Ismail et al., 2014; Seltmann et al., 2009) and seeds weight (Hufford & Hamrick, 2003; Naito et al., 2005), albinism (Takeuchi et al., 2020), as well as a poor growth and high mortality (Burgess et al., 1996; Eldridge, 1983). While ID has been much studied in short-living plants (Winn et al., 2011), often comparing fitness components between inbred and outbred plants, it is less well characterized in long-living trees (but see Griffin et al., 2019; Hedrick et al., 2016; Nickolas et al., 2019).

Most tree species avoid ID by reproducing predominantly through outcrossing (Duminil et al., 2009), a pattern also documented in tropical trees (Dick et al., 2008). In most self-compatible tropical tree species, the estimated proportion of inbred adults is near to zero even when there is a non-negligible proportion of inbred seeds (e.g. Duminil, Daïnou, et al., 2016; Duminil, Mendene Abessolo, et al., 2016). This indicates that inbred offspring are less likely to reach adult age due to their higher rate of mortality (Costa e Silva et al., 2010; Eldridge, 1983). A remarkable exception is given by *Pericopsis elata* Harms, an African timber species which was shown to produce 54% of selfed seeds in the eastern Congo Basin (Angbonda et al., 2021), questioning how this threatened species copes with ID (Angbonda et al., 2024) and what the consequences are for managing its populations. Researches on ID in trees have generally evaluated the average fitness of inbred individuals in terms of fecundity, survival and growth rate (Amorim et al., 2023; Hufford & Hamrick, 2003; Ismail et al., 2014; Seltmann et al., 2009). However, we know little about the effect of inbreeding on the functional traits and their plasticity.

Trees adapt their physiology and life cycle according to environmental conditions (Sultan, 2000). These environmental responses can be assessed using functional traits-the morphological, physiological, biochemical and phenological characteristics of plants that can be measured at the individual level (Lavorel & Garnier, 2002; Pérez-Harguindeguy et al., 2013; Violle et al., 2007; Wang et al., 2022). These traits, being related to the different functions of the plants, account for important variations in the strategy and global functioning of plants (Richardson et al., 2013) and impact fitness through their effects on plant growth, reproduction and survival (Cornelissen et al., 2003; dos Santos & Ferreira, 2020). Functional traits can vary among species, conspecific genotypes and also clonemates subject to different environments (Fyllas et al., 2009; Richardson et al., 2013). This ability of a given genotype to express different phenotypes in different environments is a phenomenon known as phenotypic plasticity (PP) (Ren et al., 2020; Sultan, 2000; Valladares et al., 2006). PP is seen as one of the main processes by which plants might acclimate to changes in environmental conditions (Gratani et al., 2000; Vitasse et al., 2010).

While the functional trait approach has often been used to compare species, it has received less attention at the intraspecific

level where the impact of ID on functional traits has rarely been investigated in tree species (Hajek et al., 2016; Violle et al., 2012). The PP of functional traits could affect fitness by allowing a genotype to better acclimate to fluctuating environmental conditions (adaptive PP). It has been hypothesized that inbreeding could reduce PP due to the reduced allelic diversity found in more homozygous genotypes (Reed et al., 2012; Swillen et al., 2015). Therefore, we hypothesize that ID could result from the expression of deleterious mutations affecting functional traits, and/ or from a reduction of adaptive PP in inbred genotypes. Studies investigating the link between PP and inbreeding or ID found various results, including a positive correlation between PP and inbreeding (e.g. Auld & Relyea, 2010; Ratz et al., 2020), no association (e.g. Buckley et al., 2019; Murren & Dudash, 2012; Sandner et al., 2022; Schlichting & Levin, 1986) or even suggested a negative correlation following the concept of developmental stability (Charlesworth & Willis, 2009).

Better understanding the interplay between ID, functional traits and PP is particularly relevant for tropical trees threatened by overexploitation. The tropical rainforests of Central Africa are experiencing unprecedented episodes of deforestation and forest degradation to sustain the livelihood of an increasing population (Aleman et al., 2018; Tyukavina et al., 2018). Logging is also affecting timber species of high commercial value on the international market, some of which suffer from a deficit of natural regeneration, like P. elata. To exploit such species in a sustainable way and avoid the depletion of natural populations, silviculture is a promising pathway but which is still little developed on native species in Central Africa (but see Kouadio & Doucet, 2009; Ouédraogo et al., 2014). The selection of good genetic material can affect the success of plantations so that it is particularly important for P. elata with a mixed mating strategy where outbred seedlings showed higher growth than inbred ones (Angbonda et al., 2024). Hence, it is crucial to analyse how inbreeding affects the PP, leaf traits and the whole-plant traits associated with growth form and competitive vigour of endemic and endangered species (Gratani, 2014).

To understand the impact of inbreeding and competition on traits associated with growth form, leaf functional traits and PP, this study focused on P. elata. Firstly, using a common garden experiment of P. elata seedlings planted at different densities and followed during 4 years (Angbonda et al., 2024; Ilunga-Mulala et al., 2021), we show that inbreeding negatively affects the whole-plant traits associated with growth and competitive vigour (diameter and maximum height), leading to ID. Then, using a set of leaf trait measurements potentially affecting photosynthesis and growth, we test two hypotheses on the possible origin of ID from the perspective of functional traits: (H1) inbreeding directly affects leaf traits; (H2) inbreeding reduces the PP of leaf traits. To test these hypotheses, we regularly measured the diameter and total height of 540 P. elata seedlings, as well as various leaf characteristics and leaf mineral content (elementome) from the base and top of the crown of 60 seedlings.

2 | MATERIALS AND METHODS

Fieldwork conducted for this study did not require any specific permits or licences. The Nelder-type experimental design was developed and established by members of the research team, who are included as co-authors of this publication.

2.1 | Species and study site

Pericopsis elata, also known as Afrormosia or Assamela, is a tall tree of high commercial value of the moist semi-deciduous African forests (Bourland, Lambert, et al., 2012). The species can reach 60 m of total height and 150 cm of diameter at breast height (Kafuti et al., 2022). It regenerates with difficulty below the crown and its natural populations are declining (Laure et al., 2014). Its natural distribution is disjoint and ranges from the lvory Coast to the Democratic Republic of Congo (Bourland, Lambert, et al., 2012). It is a light-demanding and wind-dispersed species easily recognizable by its cream or greyish scaly bark (with reddish brown spots), its compound leaves (7-11 leaflets) and its fruits (oblong indehiscent pods with 1-5 discoid seeds). The average diameter increment varies from 0.32 to 0.45 cm year⁻¹ in Cameroon and can reach 0.80 cm year⁻¹ in DRC (Bourland, Kouadio, et al., 2012; Daïnou et al., 2021). The species is listed in appendix II, CoP18#17 of CITES (https://speciesplus.net) and registered as endangered in the IUCN red list due to marked overexploitation in West Africa at the end of the 20th century (Bourland, Kouadio, et al., 2012; Hills, 2020). Silvicultural promotion of P. elata is therefore one of the best ways to support the sustainability of its populations, and to guarantee the long-term survival of the species (Ilunga-Mulala et al., 2021; Ouédraogo et al., 2014).

The plantation used for this study was installed in the suburbs of the city of Kisangani (25°15′58″ longitude East, 0°30′42″ latitude North and 495 m elevation). The seedlings for the Nelder device were obtained from seeds from 19 mother trees harvested in the Biaro forest, Tshopo province, DR Congo, where seed and pollen dispersal patterns had been studied in a 400 ha plot (Angbonda et al., 2021). Mother trees were selected based on (1) seed maturity and (2) sufficient isolation from other mother trees to ensure accuracy of seed provenance.

Seeds collected between December 2016 and January 2017 were first raised in a nursery on the campus of the Science Faculty of the University of Kisangani, planting individual seeds in 2 L plastic bags filled with forest soil. The region of Kisangani is covered with semi-deciduous forests, which are part of the Guineo-Congolian regional centre of endemism (White, 1986). The climate is of Af type, characteristic of areas without dry seasons, according to the Köppen classification. The average annual temperature is 25.1°C, and the thermal amplitude is less than 5°C. The average annual rainfall is 1672mm with a slight decrease observed between January-February and June-July (De Ridder et al., 2014). The Kisangani

region is made up of a mixture of sandstone, red clay, marl and limestone (Ilunga-Mulala et al., 2021; Mosango, 1983). Soils are Xanthic Ferralsol and Eutric Gleysols (Jones et al., 2013). The fertility is low and the cation exchange capacity (CEC) varies between 2 and 8 cmolc/kg. The soil is acidic with a pH varying between 3.5 and 5.5 (De Ridder et al., 2014).

2.2 | Sampling design, inbreeding and growth measurement

After a year in the nursery, when P. elata seedlings had reached heights of 40-160cm (median around 65cm), 648 seedlings from 19 mother trees were transplanted in October 2017 in three replicates following the la design proposed by Nelder (1962) and later applied in a context of research on monospecific tree plantations (Beeckman & Vander Mijnsbrugge, 1990). Each design occupies an area of 908 m² where seedlings are located along 12 concentric circles, each containing 18 evenly spaced saplings, ensuring a density gradient (Nelder, 1962). The saplings from the innermost and outermost circles, placed at a distance of 0.6 and 15.6 m from the centre of the circles, respectively, formed buffer zones and were not measured. Therefore, 180 out of 216 saplings were followed for each of the three replicates, totalling 540 surveyed seedlings. Local spacing between neighbouring plants ranged from 0.2×0.2 to 3.8×3.8m (Figure 1). Each of five levels of competition illustrated on Figure 1 combines two concentric circles of the Nelder device.

The data used in this study were acquired in the first 4 years after planting. In October 2017 and then every 3 months from December 2017 to December 2021, we measured the total height (*H*max) from the collar to the terminal bud of the longest branch and the diameter at 10 cm from the collar of the 540 stems. A graduated ruler and a calliper were used to measure *H*max and diameter, respectively. The Telefix dendrometer was used from 2021 for the tallest plants. *H*max and diameter are whole-plant traits associated with growth form, the species position in the vegetation's vertical light gradient, competitive vigour, etc. (Pérez-Harguindeguy et al., 2013).

We assessed whether seedlings were inbred (i.e. resulting from self-pollination) or outbred (i.e. resulting from cross-pollination) following Angbonda et al. (2021) who characterized the mating system of the source population using 11 nuclear microsatellite markers. This data set has been completed with 250 additional genotyped seedlings (Angbonda et al., 2024).

2.2.1 | Leaf traits measurement

Foliar traits were collected on 60 saplings (33 inbred, 26 outbred and one for which the inbreeding status could not be assessed). We collected leaflet samples in March 2018, 2019, 2020 and 2021 from 20 saplings per Nelder device (two individuals per circle). These saplings were taken 2 by 2 from one circle to another, alternatively along the North–South and East–West axes, due to the staggered location (Figure 1a). For each sapling, whenever possible, we sampled one leaf directly exposed to the sunlight (from the top part of the crown) and another one not directly exposed to the sunlight (from the base of the crown; Kafuti et al., 2020). Each leaf bore 6 to 10 leaflets. It was not always possible to collect both exposed and shaded leaves, especially in 2021, when some plants stayed in the shade of neighbouring plants and others had only exposed leaves. In 2019, 2020 and 2021, chlorophyll content (Chl) was measured in the field before leaflet collection using a KONICA MINOLTA SPAD-502 chlorophyllometer plus (Coste et al., 2010; Rozendaal et al., 2006; Xiong et al., 2015). The chlorophyll content of leaves is one of the most significant parameters of the physiological state of plants, enabling them to absorb solar radiation and carry out the process of photosynthesis (Silla et al., 2010).

A first set of traits included those often used to characterize the leaf economic spectrum: leaf fresh mass (LFM), leaf dry mass (LDM), leaf area (LA), specific leaf area (SLA), leaf dry matter content (LDMC), chlorophyll content (ChI), stomatal density (SD), stomatal size (SS) and maximum stomatal water vapour conductance (Gmax). These traits being involved in photosynthesis, they could express strategies of resource acquisition and phenotypic responses to environmental conditions affecting growth rate, as detailed below for each trait. They were measured each year on 60 plants, except ChI that was not measured in 2018 and stomatal traits that were collected on 30 trees in 2019 and 2020, and on 60 trees in 2021.

A second set of traits characterizing the elementome is the leaf composition in some key elements (Al, Ca, Fe, K, Mg, Mn, P and Zn), which were measured for 20 samples collected in 2020 to assess how element concentration was influenced by plant density, leaf exposure to light and inbreeding.

All leaflets collected were scanned using an HP Scanjet 5590P scanner. The high resolution images (1200 dpi) thus obtained were analysed with the software ImageJ-win64 (Schneider et al., 2012) to determine the LA (mm²). Leaflets were oven-dried at 70°C for 72 h and then stored in the envelopes with dehydrated silica gel. The dry mass of leaflets without their petioles (LDM, mg) was weighed using a precision balance (0.001 g). The SLA ($mm^2 mg^{-1}$) was calculated by dividing the leaf area by the dry mass (LA/LDM) (Pérez-Harguindeguy et al., 2013). SLA responds to a wide range of environmental stimuli (Díaz et al., 2016; Rozendaal et al., 2006; Velikova et al., 2020) and correlates with leaf chemical composition, photosynthesis rate, leaf longevity and carbon investment (Pierce et al., 2017). Leaflet dry matter content (LDMC=LDM/LFM) was also considered as it predicts leaf nitrogen content and soil fertility (Hodgson et al., 2011), and is sometimes better correlated with primary production than SLA in inter-species comparisons (Smart et al., 2017).

Stomatal characteristics of the abaxial surface of leaves were assessed in 2019 and 2020 on one or two leaflets (one exposed and one shaded) for each of 30 seedlings, and in 2021 on five or 10 leaflets (five exposed and five shaded) on each of 56 seedlings. To this end, a layer of clear nail polish was applied to the midsection of the leaflet, between the central vein and the leaflet margin. After the varnish dried overnight, the impression on the leaflet was



	Level 1	Level 2	Level 3	Level 4	Level 5
Area per plant (m ²)	0.06-0.12	0.21-0.39	0.71-1.30	2.37-4.34	7.94-14.49
Density (stems.ha-1)	690-1,260	2,304-4,219	7,702-14,079	25,794-46,613	86,044-158,75
Number of planted trees followed	108	108	108	108	108
Spacing (m)	2.8 x 2.8- 3.8 x 3.8	1.5 x 1.5 2.1 x 2.1	0.8 x 0.8 1.0 x 1.0	0.5 x 0.5 0.6 x 0.6	0.2 x 0.2 0.3 x 0.3

FIGURE 1 Nelder device. (a) Aerial view of one of the three replicates of the Nelder device. The black dots inside the competition circle represent the plants selected and monitored each year for leaf trait measurements. For each level of competition are given (b) the Hegyi's competition index (details in Ilunga-Mulala et al., 2021) and (c) the slenderness index: The ratio of total height to diameter (Kafuti et al., 2022). The slenderness index provides an expression of the social standing of trees in the Nelder device. With a declining slenderness index, the stability of trees in the system's competitive gradient rises. A slenderness value greater than or equal to 100 indicates that a tree is extremely mechanically weak. The red points in graphs (b and c) represent the average. The appended table reports information on the three Nelder replicates.

removed with tape, mounted on a glass slide and digitized using the Toupview UH-CMOS camera (Hangzhou Toup Tek Photonics Co., Ltd., Zhejiang, China) attached to the Olympus SZX-7 stereo microscope (Kafuti et al., 2020), taking five images for each leaflet. The length and width of stomata were measured on three stomata for each image, that is 15 stomata per leaflet and 7080 stomata for 472 leaflets sampled during the first 4 years of planting. Leaflets with poor quality prints were excluded from the stomatal measurements. Length (*Ls*, µm), width (*Ws*, µm) and SD (number per unit LA, mm⁻²) were determined using the ObjetJ plug-in in ImageJ-win64 (Schneider et al., 2012). The SS was obtained from the length of the

stomata: stomatal size= $0.25 \times Ls^2$ (Taylor et al., 2012). The maximum stomatal conductance to water vapour (Gmax, en mole m⁻² s⁻¹) was estimated as:

$$G \max = \frac{SD * a \max * \frac{d}{v}}{dp + \frac{\pi}{2} * \sqrt{\left(\frac{a \max}{\pi}\right)}}$$

with SD, the stomatal density, *d* is the diffusivity of water vapour to air at 25°C (=2.43 10^{-5} m²s⁻¹), *v* is the molar volume of air at 25°C (0.024 m³ mole⁻¹), *dp* is the depth of the ostiole. Since guard cells have an elliptical shape, *dp* is approximated by the stomatal width (Ws, µm)

and the maximum pore area is calculated from the stomatal length (*Ls*, μ m) using the formula of an ellipse: $amax=Ls \times Ws$ is the maximum surface of the completely open ostiole (μ m²) (Douhovnikoff et al., 2016; Kafuti et al., 2020). Note that the maximum stomatal conductance is higher (~20%) than the operating stomatal conductance (Douhovnikoff et al., 2016; Dow et al., 2014). The characteristics of stomata play a fundamental physiological role in the life of the plant (photosynthesis and transpiration) (Djinet et al., 2016). Stomatal conductance controls gas exchange between the leaves and the environment (Douhovnikoff et al., 2016; Flexas et al., 2009). This parameter is influenced by various environmental factors, such as light, temperature, humidity and water availability (Buckley, 2019).

Half of each leaflet sampled for the year 2020 in one of the replicates of the Nelder device was used for chemical analyses at the Laboratory of Plant Ecology and Biogeochemistry of the 'Université Libre de Bruxelles'. Leaflet samples were dried at 65°C and ground into a cutting mill. About 0.5 g of plant sample was dry ashed for 12 h at 450°C in a muffle furnace. Ashes were dissolved in 1mL of nitric acid (65%), heated on a hot plate for 10min. The final clear solutions were filtered using Whatman filter paper No. 41 and diluted to 15 mL. Element concentrations Al, Ca, Fe, K, Mg, Mn, P and Zn of the solutions were measured by inductively coupled plasma optical emission spectrometry (iCAP PRO, Thermo Scientific, Cambridge, United Kingdom). The precision and accuracy of the analyses were monitored using a certified reference material (Peach Leaves, NIST 1547). Analyses agreed with certified values to within \pm 5%. The nutrient content in leaflets provides information on the ability of seedlings to acquire resources (Pérez-Harguindeguy et al., 2013).

2.3 | Statistical analyses

To assess the impact of ID on the whole-plant traits associated with growth form and competitive vigour (diameter and maximum height), we modelled the maximum height or diameter growth of each plant over time using logistic regression (using R package 'stats' version 4.1.3.), separately for each competition level and inbred versus outbred plants. The growth asymptote (parameter 'a') was compared between inbred and outbred plants. The maximum height and diameter measured during the four measurement campaigns of leaf traits was also modelled using two types of linear mixed models implemented in the R package 'ImerTest' version 3.1.3 (Kuznetsova et al., 2017) and 'Ime4' version 1.1-31 (Bates et al., 2015). First, we considered all explanatory variables (year, level of competition, Nelder replicate, inbreeding and mother tree) as random factors in order to assess the proportions of variance explained by each of these factors. Second, we considered the variables year, level of competition and inbreeding as fixed factors, leaving the variables mother tree and Nelder replicate (plot) as random factors, in order to test the differences between states of the fixed factors using the R package 'emmeans' version 1.8.5 (Lenth et al., 2025).

For leaf traits, we performed two linear mixed models to assess the extent of intra- and inter-individual trait variation. The first model aimed to partition the variance of leaf traits among the different explanatory variables (year, level of competition, leaf exposure to light and inbreeding), which were all treated as random factors. In the case of elementomes, as they were measured in a single year, explanatory variables were restricted to level of competition, exposure to light and inbreeding. The second model was used to assess differences in means between states of explanatory variables of interest (levels of competition, leaf exposure to light, year and inbreeding), which were declared as fixed factors, with tree identifier and Nelder replicate used as random factors. The mother tree could not be included in these models due to the limited sample size for leaf traits (60 trees).

Phenotypic plasticity of leaf traits was quantified across years, exposure to light or the density gradient using the relative distance plasticity index (*RDPI*) ranging from 0 (no plasticity) to 1 (maximum plasticity) (Valladares et al., 2006):

$$RDPI = \frac{1}{n} \sum_{\substack{i \neq i'}}^{n} \frac{(x_{ij} - x_{i'j'})}{(x_{ij} + x_{i'j'})}$$
$$j, j'$$

where *i* represents a given level of the environmental treatment (year, levels of competition or exposure to light), *j* refers to the individual number identification, x_{ij} is the trait value of *j* in condition *i*, and the sum is taken over all possible pairs of measures taken in distinct environments for different $(j' \neq j)$ or the same (j'=j) individuals, making a total of *n* pairs. When estimating plasticity across years or exposure to light, we computed RDPI using only measures taken on the same individuals to compute RDPI across competition levels $(j' \neq j)$. RDPI was computed for each leaf trait, separately for inbred and outbred plants. To test whether RDPI was lower in inbred than outbred plants, we compared the difference with the one obtained after 1000 random permutations of the inbreeding status of seedlings. The permutation test was used to compare the plasticity of leaf traits.

To determine the link between photosynthetic capacity and growth, pairwise correlations were computed with the 'PerformanceAnalytics' version 2.0.4 R package to test the relationship between foliar traits and the height or diameter growth of inbred and outbred plants.

3 | RESULTS

The paternity analysis showed that of the 540 planted seedlings, 267 were inbred and 192 were outbred. The results were inconclusive or lacking for the remaining 81 seedlings, which were therefore not considered for statistical analyses.

3.1 | Stand competition and inbreeding effects on traits associated with growth

The total height (Hmax) and diameter were lower in inbred than in outbred plants (Figure 2, Table 1), a difference not observable until



FIGURE 2 Diameter (a) and total height (b) of P. elata seedlings as function of the age for outbred (violet) and inbred plants (green). 'a' is the estimator of the growth asymptote of the logistic model. Confidence bands were generated using bootstrapping procedure. Vertical arrows along the growth curves indicate when leaf traits measurements were taken: 150, 510, 870 and 1230 days after transplantation.

510 days after transplantation. Among the four measurement campaigns of leaf traits, the second one (2019, c. 510 days after transplantation) has recorded the highest mean growth rate in height and diameter (Figure 2).

As expected, diameter and height growth rates increased with less competition (Figure S1) and plants grew slowly after 1000 days under the highest competition levels (Figure S1A,F). According to the estimates of the asymptote (a) of the logistic models, the higher growth of outbred than inbred plants occurred at all competition levels, except at the intermediate level where a estimates had overlapping 95% confidence intervals (level 3, Figure S1C,H). The difference in growth between outbreds and inbreds tended to be highest at the lowest level of competition (Figure S1E,J).

The partition of variance indicates that the level of competition, year and Nelder replicate (plot) had the highest impact on height and diameter growth (Figure 3c), often explaining 19% to 25% of the variation, while inbreeding and mother tree explained 4% to 5% of the variation.

3.2 Determinants of leaf traits variation

Plant leaf characteristics varied substantially as a function of year, planting density, and/or leaf exposure to light (Figure 3; Table 1). On the contrary, inbreeding had no significant impact on any of the studied leaf traits (Table 1), despite the fact that about 4% to 5% of the variation in plant growth was explained by inbreeding and by the mother tree (Figure 3c).

Competition affected in particular SLA, explaining 11.2% of its variance, Hmax (24.5%), Diameter (19.5%) and iron concentration (Fe, 60.3%) (Figure 3a-c). SLA increased by about 50% from lowest to highest competition, an effect due to a decrease in leaf mass rather than a change in LA (Table 1). Similarly, leaf Fe concentration increased about twofold along the competition gradient (Table 1).

Year affected most of the leaf traits (e.g. 38% explained variance for dry mass content) but had a non-significant impact on stomata size or density (Table 1). There was no regular trend with year, as it was most often the Year 2019 that stood out from the other ones (higher leaf mass, dry mass content, LA, chlorophyll content and Gmax, lower SLA; Table 1; Figure S2).

Leaf exposure to light affected mostly SD (13.3%, albeit nonsignificantly), fresh leaf mass (higher for exposed leaflets) and all elements (28.7% to 69.8%) except Fe concentration. Leaflets well exposed to sunlight had lower concentration in Al, Ca, Mg and Mn, but higher concentration in K, P and Zn (Table 1).

Phenotypic plasticity of inbred and 3.3 outbred plants

Overall, PP was moderate for all leaf traits, with RDPI in the range 0.15-0.32 for most traits (Figure 4) but lower plasticity was found for chlorophyll content (CHL: 0.11), stomata size (SS: 0.04 to 0.06) and stomatal conductivity (Gmax: 0.04 to 0.05). Plasticity was nearly always slightly lower across exposure to sunlight (measured within year and tree) than across years (measured within tree and sun

TABLE 1	Means and st	andard errors	of growth and	foliar traits acr	oss years, con	petition level	s, position of th	ne leaf within t	he crown and i	nbreeding.			
	Year				Competition					Exposure to su	ц	Inbreeding	
Traits	2018 [1.42 year]	2019 [2.42 year]	2020 [3.42 year]	2021 [4.42 year]	Level 1 [very low]	Level 2 [low]	Level 3 [medium]	Level 4 [high]	Level 5 [very high]	Shaded	Exposed	Inbred	Outbred
Growth traits													
Diameter	I	32±1.47a	43.10±1.47b	$53.10 \pm 1.47c$	$25.80\pm1.96a$	$37.90 \pm 1.93b$	$43.90 \pm 1.94b$	$52.3 \pm 1.94c$	53.80±1.74c	Ι	Ι	39±1.62a	46.4±1.55b
Height	I	263±9.84a	$342 \pm 9.84b$	402±9.84c	275 ± 13a	$360 \pm 12.90b$	$357 \pm 13b$	$351\pm13b$	$335 \pm 11.70b$	I	I	316±10.80a	$355 \pm 10.30b$
Leaf structura	I traits												
LFM	0.29 a±0.01	$0.34 \pm 0.01b$	0.26±0.01a	$0.24 \pm 0.01a$	0.34±0.01a	0.30±0.01ab	$0.28 \pm 0.01b$	$0.25 \pm 0.02b$	$0.24 \pm 0.01b$	0.25±0.01a	$0.31 \pm 0.01b$	0.27 ± 0.01	0.29 ± 0.01
LDM	0.08±0.01a	$0.14 \pm 0.01b$	$0.07 \pm 0.01a$	0.06±0.01a	$0.11 \pm 0.01a$	0.09±0.01ab	$0.08 \pm 0.01b$	0.08±0.01ab	$0.07 \pm 0.01 b$	0.08 ± 0.01	0.09 ± 0.01	0.08 ± 0.01	0.09 ± 0.01
LDMC	0.28±0.02a	0.41 ±0.02b	0.28±0.01a	0.29±0.02a	0.33 ± 0.02	0.31 ± 0.02	0.30 ± 0.02	0.33 ± 0.02	0.31 ± 0.02	0.33 ± 0.02	0.30 ± 0.02	0.31 ± 0.02	0.31 ± 0.02
ΓA	$17.20 \pm 0.83a$	$21.30 \pm 0.82b$	15.00±0.86a	18.40±0.96ab	18.10 ± 1.01	17.70 ± 1.02	16.90 ± 1.04	17.90 ± 1.55	19.20 ± 0.90	17.40 ± 0.72	18.60 ± 0.72	17.40 ± 0.81	18.50 ± 0.76
SLA	$22.50 \pm 1.70a$	15.60±1.69ab	24.60±1.72ab	$32.6 \pm 1.79b$	18.10±1.92a	21.08±1.93ab	23.70±1.95ab	25.80±2.55ab	29.90±1.80b	25.90 ± 1.62	21.80 ± 1.61	23.6 ± 1.66	24.40 ± 1.71
CHL	I	35.80±0.86a	27.20±0.90b	$28.60 \pm 0.99b$	$34.20 \pm 1.07a$	$32.10 \pm 1.04 ab$	$28.60 \pm 1.1b$	$28.00 \pm 1.65b$	$30.10\pm0.96ab$	30.70 ± 0.82	30.80 ± 0.81	30.9 ± 0.81	30.5 ± 0.87
SS	I	97.70 ± 2.86	91.40 ± 2.88	92.50 ± 2.61	96.90 ± 3.17	91.7 ± 3.43	92.20 ± 3.01	94.00 ± 3.72	94.70 ± 3.58	95.10 ± 2.68	92.70±2.69	93.6±2.80	94.2 ± 2.90
SD	I	246 ± 16.30	264 ± 16.30	222 ± 12.20	268 ± 17	268 ± 21.30	221 ± 14.50	237 ± 21.10	227 ± 17.40	218 ± 12.70	259 ± 12.80	237 ± 13.40	240 ± 14.10
Gmax	I	1.74±0.02a	$1.71 \pm 0.02ab$	$1.61 \pm 0.01b$	1.75 ± 0.02	1.69 ± 0.02	1.67 ± 0.03	1.65 ± 0.03	1.65 ± 0.03	1.67 ± 0.01	1.70 ± 0.01	1.68 ± 0.01	1.68 ± 0.01
Leaf chemical	traits												
AI	I	l	117.61 ± 55.41	I	88.20±13a	86±14a	127.50±18ab	$124.70\pm14ab$	$164\pm13.50\mathrm{b}$	$147\pm9.34a$	$88.40 \pm 8.94b$	110 ± 10.81	126 ± 8.66
Ca	I	I	5342 ± 3598	I	6307 ± 882	6026 ± 913	4922 ± 1226	4258 ± 913	5458 ± 882	8022±608a	2767 ± 582b	5536 ± 704	5253 ± 564
Fe	I	I	58.45 ± 27.01	I	36.50±6.44a	40.20±6.67ab	55.40±8.85ab	66.50±6.67ab	94.30±7.6b	63±4.20	54.20 ± 4.03	54.40 ± 5	62.80 ± 4.11
¥	I	I	4211 ± 2141	I	4346±636	5118 ± 659	4114 ± 885	3462 ± 659	4230 ± 637	3078±439a	$5430 \pm 421b$	4887 ± 508	3622 ± 407
Mg	I	I	1188 ± 522	I	990 ± 158	1588 ± 164	1091 ± 220	1074 ± 164	1256 ± 158	1424 ± 109	9.76 ± 105	1302 ± 126	1098 ± 101
Mn	I	l	13.49 ± 7.17	I	16.30 ± 1	13.80 ± 1	12.60 ± 2	12.40 ± 1	11.50 ± 1	$17.90 \pm 1a$	$8.80 \pm 1b$	11.60 ± 1	15.10 ± 1
д	I	I	1229 ± 392	I	1167 ± 110	1379 ± 113	1230 ± 152	1222 ± 113	1161 ± 110	980±75a	$1484\pm72b$	1302 ± 87	1161 ± 70
Zn	I	I	17.01 ± 7.16	I	16.80 ± 2.66	18.60 ± 2.76	13.60 ± 3.62	18.70 ± 2.76	15.90 ± 2.66	13±1.67a	$20\pm1.60b$	16.70 ± 2.12	16.70 ± 1.70
<i>Note</i> : The ag	e mentioned h∈	starts from t	the germination	in the nursery i	until the date o	f measurement	. Values in black	k indicates that	significant diffe	srences occur	according to f	actor of intere	st, while

values in grey indicate no significant differences. The letters (a-c) indicates which means are significantly different at an α =0.05. All chemical traits are expressed in µg/g; leaf trait units are provided in Methodology Section 2.2. N

BRITISH ECOLOGICAL Journal of Ecology





FIGURE 3 Variance partitioning of (a) nine leaf morphological and physiological traits (CHL, chlorophyll content; Gmax, maximum stomatal conductance; LA, leaf area; LDM, leaf dry mass; LDMC, leaf dry matter content; LFM, leaf fresh mass; SD, stomatal density; SLA, specific leaf area; SS, stomatal size), (b) leaf concentration of eight elements and (c) two traits associated with growth (stem diameter and height) across five explanatory categorical variables expressing the effect of age (Year, except for elements which were measured on a single year), light exposure (leaves from top or bottom of the crown, except for height and diameter), competition (five levels of stand density), inbreeding (inbred versus outbred plants) and mother tree (only for height and diameter). The residual variation (Error) includes unexplained variation within and between individuals. For each level where the explained variance is higher than 10%, the value is printed on the corresponding stacked bar.

exposure level) or competition levels (measured within year and sun exposure level). Plasticity across years and competition levels was similar, except for SLA showing higher plasticity across competition levels (0.25) than across years (0.21) and across exposure to sunlight (0.16).

Permutation tests found higher plasticity in outbred than inbred plants (one-sided tests) only for chlorophyll content (CHL) across years (p=0.029) and competition levels (p=0.032), for fresh leaf mass (LFM) across exposure to sunlight (p=0.020) and competition levels (p=0.062, marginal significance), and for stomatal conductivity

(Gmax) across competition levels (p=0.072, marginal significance). The sole case of a statistically significant reverse pattern-higher plasticity for inbred than outbred plants-was found for leaflet dry mass content (LDMC) across competition levels (p=0.028).

3.4 | Relationship between leaf traits and growth

Correlations between leaf traits for inbred and outbred plants are presented in Table S1. Growth in diameter was negatively related to



FIGURE 4 Phenotypic plasticity of leaf traits estimated as the relative distance plasticity index (*RDPI*) and its 95% confidence interval, according to three environmental gradients: (a) between upper and lower leaf position in the crown (sunlight gradient), (b) across year (ontogenetic plasticity) and (c) across different levels of competition (environmental plasticity). For each trait, the average *RDPI* is compared between inbred (green) and outbred (purple) individuals, and tested following a permutation test. *: significant difference (p < 0.05).

SLA, both in inbred plants (R^2 =0.49, p < 0.001) and in outbred plants (R^2 =0.22, p < 0.01) (Figure 5). Growth in height was significantly related to SLA in inbred plants (R^2 =0.32, p < 0.01) but not in outbred plants (R^2 =0.017, p>0.05).

4 | DISCUSSION

This study shows that inbreeding and competition (planting density) have a significant impact on the height and diameter growth of 1- to 4-year-old *P. elata* saplings. While leaf traits varied with competition, age and/or leaf exposure to direct sunlight, they were not affected by inbreeding. However, the PP of some leaf traits was slightly lower



FIGURE 5 (a) Diameter ($a_{Diameter}$) and (b) height (a_{Height}) growth estimates of *Pericopsis elata* as function of the specific leaf area (SLA) for outbred (violet) and inbred (green) saplings. Regression line, coefficient of determination (R^2) and statistical significance: *** for p < 0.001; ** for p < 0.01; ns for non-significant. Horizontal and vertical bars indicate the standard errors of means.

in inbred than outbred saplings. On the contrary, the SLA of inbred saplings was more correlated with height and diameter growth than that of outbred saplings. The various effects observed are discussed below.

4.1 | Higher growth performance of outbred saplings

The significant impact of competition on sapling growth has been reported by other authors in a natural environment (Rozendaal et al., 2020) or in plantations (Angbonda et al., 2024; Ilunga-Mulala et al., 2021). These results underline the relevance of density control in forest plantations (Pommerening & Grabarnik, 2019). High planting density affects the diameter growth more than the height growth (Figure S1) so that slenderness increases with planting density (Figure 1c). This confirms an important theory in dendrometry, according to which tree species generally respond to light competition by investing more energy in height growth than in diameter growth (Brunner & Nigh, 2000).

After 53 months in plantation, outbred P. elata plants showed higher growth performance in height and diameter than inbred plants for all levels of competition, a result also reported in Angbonda et al. (2024). Similar results were reported in Prunus dulcis (Oukabli et al., 2001) and Shorea laxa seedlings (Takeuchi et al., 2020). All these studies show that outbred plants outperform inbred plants, confirming a strong link between inbreeding and growth (Burgess et al., 1996). Furthermore, our results showed no significant difference between inbred and outbred plants until around 510 days after transplantation. These results suggest that the deleterious effects of ID are only observable around a year and a half after transplanting P. elata seedlings. Similar observations were reported by Stacy (2001) on the growth of Syzygium rubicundum and Shorea cordifolia, where no effect of the cross or parent tree was observed on growth in the first year after sowing. Hardner and Potts (1995) also reported a trend for ID to increase with age in Eucalyptus globulus seedlings. In Eucalyptus regnans plants, the growth of inbred and outbred did not differ significantly up to the age of 2 years, with very high mortality in inbred families from the age of 13 (Eldridge, 1983). The effects of inbreeding are most apparent at the seed formation stage, and less at the seed germination and early seedling growth stages (Stacy, 2001). This phenomenon was not tested in P. elata because it was not possible to follow the fate of inbred versus outbred ovules before seed dispersal. In view of these observations, we confirm that inbreeding negatively affects the height and diameter growth of young P. elata plants. Furthermore, the impact of inbreeding increases with seedling age.

4.2 | Leaf traits determined by environmental conditions but not by inbreeding

We found considerable variation in chemical and foliar traits of *P. elata* trees that were mostly explained by plant development stage (year), light exposure and the level of competition. By contrast, inbreeding had no measurable impact on these leaf traits, refuting our hypothesis H1 that inbreeding negatively affects leaf traits.

We must remain cautious with the later interpretation because while growth was measured on more than 500 plants, leaf traits were measured on 60 plants (20 for leaf elementome), limiting the statistical power to detect ID. The acquisition of functional traits is time consuming and costly, so we opted firstly to take repeated measurements on a limited number of plants to be able to assess the sources of PP (i.e. variation within each genotype). Nevertheless, our data are conclusive enough to affirm that inbreeding has a negligible Journal of Ecology

impact on leaf trait values compared with the impact of year, compe-

For most leaf traits, notably LDM, LDMC, LA, SLA, CHL and Gmax, the first source of explained variation is year, where measurements made in 2019, about 510 days after transplantation, often departed from the other ones (Table 1; Figure S2). This is concomitant with the highest growth rates recorded (Figure 2). These results are in line with previous studies reporting that the characteristics of the leaves and/or the whole plant change with age (e.g. Laforest-Lapointe et al., 2014), including leaf chlorophyll content (Wang et al., 2020), and independently of the competitive and light environment (Coleman et al., 1994).

tition or light exposure (Figure 3).

Some leaf traits responded to the competition gradient. With increasing plant density, we observe in particular a decrease in leaf dry and fresh mass (LDM and LFM), increasing SLA, a decrease in chlorophyll concentration (CHL) and an increase in Fe concentration, the sole element responding to the density gradient. These variations are explained by the fact that different levels of competition create different microenvironments (Ren et al., 2020). According to Ackerly et al. (2002), low SLA contributes to leaf survival and nutrient retention, the situation observed under low competition. Silvicultural practices therefore affect the mineral retention capacity of *P. elata* saplings (Ebuy et al., 2016).

Exposure to sunlight had little impact on most leaf traits, except that exposed leaves had higher fresh mass (LFM). By contrast, sun exposure had a very strong impact on the concentration of most leaf elements, except Fe, and in different directions according to the specific element considered. More specifically, Al, Ca, Mg and Mn were more concentrated in the shaded foliage at the base of the crown, while K. P and Zn were more concentrated in the wellexposed foliage at the top of the crown. A similar observation was reported by Hedde et al. (2007), where leaves exposed to full sunlight had lower nutrient levels and a more capacitive antioxidant system than leaves in the shade. The abundance of these three elements (K, P and Zn) in leaves fully exposed to sunlight is explained by their relevance to primary growth, given that potassium controls water pressure (opening and closing of stomata) (Santiago et al., 2012), phosphorus and zinc improve photosynthetic pigment production and growth (Richardson & Simpson, 2011; Walker et al., 2014; Zhang et al., 2023). Furthermore, Fyllas et al. (2009) show that mean annual temperature is negatively correlated with leaf concentrations of nitrogen, phosphorus and potassium. This might also explain why leaves exposed to sunlight contain more nutrients for photosynthetic growth than leaves located at the base of the canopy. Soil fertility plays a major role in determining the chemical properties of the leaves of specific tree species, in particular their N, P, K, Mg, Cu and Al content (Fyllas et al., 2009; Heineman et al., 2016). However, in the relatively homogeneous pedological environment where the Nelder devices were installed (unpublished results), the variability of the leaf chemical traits does not appear correlated with local soil variability and would depend primarily on the strategy of capture and conservation of

Journal of Ecology

resources affected by the position of the leaves in the crown and access to light.

In contrast to the results of Aranda et al. (2001), where the SD of *Pinus sylvestris* is positively correlated to light exposure, we found no impact of sunlight exposure, competition and seedling age on SD or size (SS and SD; Table 1). Only Gmax, which depends on SD and size, showed significant variation according to age. However, tree identity explained a large portion of the variation in SD and size, both for inbred and outbred seedlings (Figure S3), suggesting that these traits have a strong genetic determinism (i.e. high broad-sense heritability). In Amazonia, Camargo and Marenco (2011) have suggested that stomatal variation depends on the genetic background of each species. Our results suggest furthermore that the importance of the genetic background also applies at the intraspecific level.

4.3 | Phenotypic plasticity of some leaf traits is slightly higher in outbred plants

For most traits and environmental contrasts, PP did not differ significantly between inbred and outbred plants (Figure 4) and there was no dominant trend either. However, we found that PP was slightly but significantly higher for outbred than inbred plants for CHL across years and the competition gradient, for Gmax across the competition gradient, and for LFM across the light exposure and competition gradients (but LDMC showed higher plasticity for inbreds across the competition gradient). Hence, we find only moderate support for hypothesis H2 stating that inbreeding reduces PP.

Auld and Relyea (2010) report an effect of inbreeding on PP while Murren and Dudash (2012) found no significant differences between inbred and outbred plants. It is usually argued that plants with greater morphological plasticity are better able to adapt to heterogeneous environments (Abdusalam & Li, 2018), possibly contributing to ID if inbred plants showed lower plasticity due to a reduction of their allelic diversity at functional genes. Our results indicate that the higher plasticity of outbred *P. elata* plants for some foliar characteristics, particularly chlorophyll content, maximum water vapour conductance and fresh mass, might confer a higher potential for acclimation in a changing environment. We can, however, not demonstrate that such an effect is responsible for the ID detected on *H*max and diameter.

For tree species to survive rapid anthropogenic climate change, they must adapt their phenotypes through plasticity (Nicotra et al., 2010) by adjusting trait values to suit specific conditions (Anderson et al., 2012). Further work on the PP of *P. elata* and other tree species is still needed to better understand how ongoing environmental changes will affect tree populations.

4.4 | Relationship between growth and foliar traits

In this study, we found significantly negative correlations between SLA and both maximum height growth ($R^2=0.32^{**}$) and diameter

growth ($R^2 = 0.49^{***}$) in inbred seedlings, and weaker relationships in outbred seedlings (Figure 5). These results suggest that SLA may be a better indicator of growth in inbred plants than in outbred plants. These relationships are opposite to those typically found at the interspecific level, where growth rate and SLA are positively correlated (Poorter et al., 2008; Poorter & Bongers, 2006). SLA is an indicator of the efficiency of investment in biomass for light interception. When evaluating the field performance of seedlings of 53 Neotropical tree species, Poorter and Bongers (2006) found that SLA was a reasonable predictor of interspecific variation in height growth ($R^2 = 0.18$) and interpreted their results as an adaptive consequence of the growthsurvival trade-off. The negative relationship between SLA and growth reported in our study possibly originates from the fact that we consider the growth-SLA relationship at the intraspecific level and that our measures were taken along a steep gradient of competition. High competition reduces the growth rate of diameter, due to resource limitation, but the latter could also limit the development of thick leaves, resulting in high SLA. Hence, we suggest that the negative growth-SLA correlation found in P. elata would result from environmental constraints rather than reflect an adaptive strategy.

The fact that this negative relationship manifests more strongly in inbred seedlings than outbred ones (Figure 5) is more difficult to interpret. It may reflect a higher capacity of outbred plants to compensate for environmental constraints, reflecting higher plasticity, but our plasticity measure did not detect a difference in plasticity between inbred and outbred plants for SLA, and our sample size is too small to demonstrate that the growth–SLA correlation truly differs according to inbreeding.

5 | CONCLUSION

The aim of this study was to evaluate the impact of inbreeding and planting density on the whole-plant traits associated with growth form and competitive vigour, the variability of leaf functional traits and their PP, using young individuals of P. elata. We found lower height and diameter for inbred seedlings compared with outbred seedlings. This ID was not clearly reflected in the values of leaf traits. However, some leaf traits had slightly higher PP in outbred seedlings. Whether this difference in plasticity could explain, or at least contribute, to ID remains unclear. Our results highlight the importance of taking into account the selfed/outcrossed origin of P. elata seeds and density control when establishing sustainable forest plantations in a context of environmental change. The implications for management and conservation of P. elata are discussed by Angbonda et al. (2024). Few studies have investigated so far the potential interplay between ID, functional traits and PP at the intraspecific level. We argue that such studies should be replicated in different biological models to assess whether general patterns emerge.

AUTHOR CONTRIBUTIONS

Hans Beeckman and Nils Bourland conceived and implemented the plantations under Nelder designs. Dieu-Merci Assumani Angbonda

Journal of Ecology

collected the seeds in the Biaro forest, set up and monitored the saplings in the nursery under Nils Bourland's and Olivier J. Hardy's supervision. Jean Pierre Ngongo, Crispin Ilunga-Mulala, Espoir Ibrahim and Hulda Hatakiwe collected data in the field. Jean Pierre Ngongo and Espoir Ibrahim carried out laboratory treatment of leaf traits. Dieu-Merci Assumani Angbonda and Olivier J. Hardy genotyped the seedlings to identify the ones resulting from selfing. Jean Pierre Ngongo analysed the phenotypic data with the support of Olivier J. Hardy, Chadrack Kafuti, Nils Bourland, Hans Beeckman, Thomas Drouet, Léopold Ndjele and Yves Brostaux. Jean Pierre Ngongo wrote the manuscript under the supervision of Olivier J. Hardy. The final draft of the manuscript received contributions from all authors. All authors have read and agreed to the published version of the manuscript.

ACKNOWLEDGEMENTS

We thank the Royal Museum for Central Africa for installing the Nelder device used in this study to monitor the growth of Pericopsis elata. We also thank the team of the 'LECAFOR' Ecology and Forest Management Laboratory for collecting seeds and taking care of the nursery work. We express our gratitude to the Yangambi Wood Biology Laboratory, Belgian Directorate-General for Development Cooperation and Humanitarian Aid (DGD) via PilotMAB and European Union (EU) via the Center for International Forestry Research (CIFOR); FORETS project (EEC40) to facilitate leaf trait analysis. We thank Saskia Sergeant and Ecren Ariöz for their help in genotyping samples. We thank the CFT 'Compagnie Forestière de Transformation', especially to its Managing Director, K. Ammacha, for welcoming us during our fieldwork and providing invaluable assistance with sample collection. We are particularly grateful to ARES 'Académie de Recherche et d'Enseignement Supérieur' for granting Jean Pierre Ngongo a PhD scholarship, which enabled him to collect data in the field. Molecular laboratory analyses were funded by Belgian Fund for Scientific Research (F.R.S-FNRS). Finally, we thank two anonymous referees who provided insightful comments to improve our work.

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests.

PEER REVIEW

The peer review history for this article is available at https://www. webofscience.com/api/gateway/wos/peer-review/10.1111/1365-2745.70082.

DATA AVAILABILITY STATEMENT

Data used in this study were collected between 2017 and 2021 and can be found at https://doi.org/10.5281/zenodo.13927593 (Ngongo, 2024).

ORCID

Jean Pierre Ngongo 🕩 https://orcid.org/0009-0006-1383-5009 Crispin Ilunga-Mulala 🕩 https://orcid.org/0000-0003-4456-0846

REFERENCES

- Abdusalam, A., & Li, Q. (2018). Morphological plasticity and adaptation level of distylous Primula nivalis in a heterogeneous alpine environment. *Plant Diversity*, 40, 284–291. https://doi.org/10.1016/j.pld. 2018.11.003
- Ackerly, D. D., Knight, C. A., Weiss, S. B., & Barton, K. (2002). Leaf size, specific leaf area and microhabitat distribution of chaparral woody plants: Contrasting patterns in species level and community level analyses. *Oecologia*, 130, 449–457. https://doi.org/10.1007/s0044 20100805
- Aleman, J. C., Jarzyna, M. A., & Staver, A. C. (2018). Forest extent and deforestation in tropical Africa since 1900. Nature Ecology & Evolution, 2, 26–33. https://doi.org/10.1038/s41559-017-0406-1
- Amorim, G. T. d. S., Pinto, T. d. O., Silveira, T. d. O., de Souza, M. A. A., Menezes, B. R. d. S., & Junior, P. C. D. (2023). *Catharanthus roseus*[L.] G. Don: Allogamy as the main reproductive strategy and autogamy as a supposed reproductive guarantee mechanism. *South African Journal of Botany*, 154, 32–40. https://doi.org/10.1016/j. sajb.2023.01.004
- Anderson, J. T., Panetta, A. M., & Mitchell-Olds, T. (2012). Evolutionary and ecological responses to anthropogenic climate change. *Plant Physiology*, 160, 1728–1740. https://doi.org/10.1104/pp.112. 206219
- Angbonda, D.-M. A., Ilunga-Mulala, C. M., Bourland, N., Beeckman, H., Hatakiwe, H., Ngongo, J. P., & Hardy, O. J. (2024). Inbreeding depression affects the growth of seedlings of an African timber species with a mixed mating reproductive system, *Pericopsis elata*. *Heredity*, 133, 238-248. https://doi.org/10.1038/s41437-024-00709-x
- Angbonda, D. M. A., Monthe, F. K., Bourland, N., Boyemba, F., & Hardy, O. J. (2021). Seed and pollen dispersal and fine-scale spatial genetic structure of a threatened tree species: *Pericopsis elata* (HARMS) Meeuwen (Fabaceae). *Tree Genetics & Genomes*, 17, 27. https://doi. org/10.1007/s11295-021-01509-8
- Aranda, I., Gil, L., & Pardos, J. (2001). Effects of thinning in a Pinus sylvestris L. stand on foliar water relations of Fagus sylvatica L. seedlings planted within the pinewood. Trees, 15, 358–364. https://doi.org/ 10.1007/s004680100109
- Auld, J. R., & Relyea, R. A. (2010). Inbreeding depression in adaptive plasticity under predation risk in a freshwater snail. *Biology Letters*, 6, 222–224. https://doi.org/10.1098/rsbl.2009.0726
- Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48. https://doi.org/10.18637/jss.v067.i01
- Beeckman, H., & Vander Mijnsbrugge, K. (1990). The evolutionary learning algorithm in processing of ecological data. Case-study: Poplar tree-ring analysis. *Silva Gandav*, *55*, 87–103. https://doi.org/10. 21825/sg.v55i0.903
- Bourland, N., Kouadio, Y. L., Fétéké, F., & Lejeune, P. (2012). Ecology and management of *Pericopsis elata* (Harms) Meeuwen (Fabaceae) populations: A review. *Biotechnology*, *Agronomy*, *Society and Environment*, 16, 486–498.
- Bourland, N., Lambert, K. Y., Lejeune, P., Bonaventure, S., Julien, P., Daïnou, K., Fétéké, F., & Doucet, J.-L. (2012). Ecology of *Pericopsis elata* (Fabaceae), an endangered timber species in southeastern Cameroon. *Biotropica*, 44, 840–847. https://doi.org/10.1111/j. 1744-7429.2012.00874.x
- Brunner, A., & Nigh, G. (2000). Light absorption and bole volume growth of individual Douglas-fir trees. *Tree Physiology*, 20, 323–332. https://doi.org/10.1093/treephys/20.5-6.323
- Buckley, J., Daly, R., Cobbold, C. A., Burgess, K., & Mable, B. K. (2019). Changing environments and genetic variation: Natural variation in inbreeding does not compromise short-term physiological responses. Proceedings of the Royal Society B: Biological Sciences, 286, 20192109. https://doi.org/10.1098/rspb.2019.2109

- Buckley, T. N. (2019). How do stomata respond to water status? The New Phytologist, 224, 21–36. https://doi.org/10.1111/nph.15899
- Burgess, I. P., Williams, E. R., Bell, J. C., Harwood, C. E., & Owen, J. V. (1996). The effect of outcrossing rate on the growth of selected families of *Eucalyptus grandis*. *Silvae Genetica*, 45, 97–100.
- Camargo, M. A. B., & Marenco, R. A. (2011). Densidade, tamanho e distribuição estomática em 35 espécies de árvores na Amazônia central. Acta Amazonica, 41, 205–212. https://doi.org/10.1590/S0044 -59672011000200004
- Charlesworth, D., & Willis, J. H. (2009). The genetics of inbreeding depression. Nature Reviews. Genetics, 10, 783–796. https://doi.org/10. 1038/nrg2664
- Coleman, J., McConnaughay, M. K. D., & Ackerly, D. D. (1994). Interpreting phenotypic variation in plants. *Trends in Ecology & Evolution*, 9, 187–191.
- Cornelissen, J. H. C., Lavorel, S., Garnier, E., Díaz, S., Buchmann, N., Gurvich, D. E., Reich, P. B., Ter Steege, H., Morgan, H. D., Van Der Heijden, M. G. A., Pausas, J. G., & Poorter, H. (2003). A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian Journal of Botany*, *51*, 335–380. https://doi.org/10.1071/BT02124
- Costa e Silva, J., Hardner, C., Tilyard, P., Pires, A. M., & Potts, B. M. (2010). Effects of inbreeding on population mean performance and observational variances in *Eucalyptus globulus*. *Annals of Forest Science*, 67, 605. https://doi.org/10.1051/forest/2010018
- Coste, S., Baraloto, C., Leroy, C., Marcon, É., Renaud, A., Richardson, A. D., Roggy, J. C., Schimann, H., Uddling, J., & Hérault, B. (2010). Assessing foliar chlorophyll contents with the SPAD-502 chlorophyll meter: A calibration test with thirteen tree species of tropical rainforest in French Guiana. *Annals of Forest Science*, 67, 607. https://doi.org/10.1051/forest/2010020
- Daïnou, K., Tosso, F., Bracke, C., Bourland, N., Forni, E., Hubert, D., Kankolongo, A., Loumeto, J., Louppe, D., Ngomanda, A., Ngomin, A., Tite, V., & Doucet, J.-L. (2021). Guide Pratique des Plantations d'Arbres des forêts denses humides d'Afrique.
- Darwin, C. R. (1876). The effects of cross and self fertilisation in the vegetable. John Murray.
- De Ridder, M., Toirambe, B., Van den Bulcke, J., Bourland, N., Van Acker, J., & Beeckman, H. (2014). Dendrochronological potential in a semi-deciduous rainforest: The case of *Pericopsis elata* in central Africa. *Forests*, *5*, 3087-3106. https://doi.org/10.3390/ f5123087
- Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice, I., Garnier, E., Bönisch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T., Dickie, J., Gillison, A. N., Zanne, A. E., ... Gorné, L. D. (2016). The global spectrum of plant form and function. *Nature*, *529*, 167–171. https://doi. org/10.1038/nature16489
- Dick, C. W., Hardy, O. J., Jones, F. A., & Petit, R. J. (2008). Spatial Scales of pollen and seed-mediated gene flow in tropical rain Forest trees. *Tropical Plant Biology*, 1, 20–33. https://doi.org/10.1007/s1204 2-007-9006-6
- Djinet, A. I., Bell, J. M., Nana, R., Nguinambaye, M. M., & Tamini, Z. (2016). Évaluation des caractéristiques des stomates chez le palmier à huile (*Elaeis guineensis* Jacq.). *Journal of Applied Biosciences*, 104, 9904. https://doi.org/10.4314/jab.v104i1.2
- dos Santos, V. A. H. F., & Ferreira, M. J. (2020). Are photosynthetic leaf traits related to the first-year growth of tropical tree seedlings? A light-induced plasticity test in a secondary forest enrichment planting. *Forest Ecology and Management*, 460, 117900. https://doi.org/ 10.1016/j.foreco.2020.117900
- Douhovnikoff, V., Taylor, S. H., Hazelton, E. L. G., Smith, C. M., & O'Brien, J. (2016). Maximal stomatal conductance to water and plasticity in stomatal traits differ between native and invasive introduced lineages of *Phragmites australis* in North America. *AoB Plants*, *8*, 1-11. https://doi.org/10.1093/aobpla/plw006

- Dow, G. J., Bergmann, D. C., & Berry, J. A. (2014). An integrated model of stomatal development and leaf physiology. *The New Phytologist*, 201, 1218–1226. https://doi.org/10.1111/nph.12608
- Duminil, J., Daïnou, K., Kaviriri, D. K., Gillet, P., Loo, J., Doucet, J. L., & Hardy, O. J. (2016). Relationships between population density, finescale genetic structure, mating system and pollen dispersal in a timber tree from African rainforests. *Heredity (Edinb)*, 116, 295–303. https://doi.org/10.1038/hdy.2015.101
- Duminil, J., Hardy, O. J., & Petit, R. J. (2009). Plant traits correlated with generation time directly affect inbreeding depression and mating system and indirectly genetic structure. BMC Evolutionary Biology, 9, 1–14. https://doi.org/10.1186/1471-2148-9-177
- Duminil, J., Mendene Abessolo, D. T., Ndiade Bourobou, D., Doucet, J. L., Loo, J., & Hardy, O. J. (2016). High selfing rate, limited pollen dispersal and inbreeding depression in the emblematic African rain forest tree Baillonella toxisperma—Management implications. Forest Ecology and Management, 379, 20–29. https://doi.org/10.1016/j. foreco.2016.08.003
- Ebuy, J., Mate, J. P., Mukandama, J. P., & Ponette, Q. (2016). Chute des litières et fertilité des sols sous plantations forestières dans le bassin du Congo: Cas de la station I.N.E.R.A/Yangambi en R.D.C. *Journal of Animal & Plant Sciences*, 31, 4843–4861.
- Eldridge, K. G. (1983). Selfing effets in Eucalyptus regnans. Silvae Genetica, 32, 5–6.
- Flexas, J., Barón, M., Bota, J., Ducruet, J. M., Gallé, A., Galmés, J., Jiménez, M., Pou, A., Ribas-Carbó, M., Sajnani, C., Tomàs, M., & Medrano, H. (2009). Photosynthesis limitations during water stress acclimation and recovery in the drought-adapted Vitis hybrid Richter-110 (V. berlandieri x V. rupestris). Journal of Experimental Botany, 60, 2361– 2377. https://doi.org/10.1093/jxb/erp069
- Fyllas, N. M., Patino, S., Baker, T. R., Bielefeld Nardoto, G., Martinelli, L. A., Quesada, C. A., Paiva, R., Schwarz, M., Horna, V., Mercado, L. M., Santos, A., Arroyo, L., Jiměnez, E. M., Luizao, F. J., Neill, D. A., Silva, N., Prieto, A., Rudas, A., Silviera, M., ... Lloyd, J. (2009). Basinwide variations in foliar properties of Amazonian forest: Phylogeny, soils and climate. *Biogeosciences*, *6*, 2677–2708. https://doi.org/10. 5194/bg-6-2677-2009
- Gratani, L. (2014). Plant phenotypic plasticity in response to environmental factors. Advances in Botany, 2014, 1–17. https://doi.org/10. 1155/2014/208747
- Gratani, L., Pesoli, P., Crescente, M. F., Aichner, K., & Larcher, W. (2000). Photosynthesis as a temperature indicator in *Quercus ilex* L. *Global* and Planetary Change, 24, 153–163. https://doi.org/10.1016/S0921 -8181(99)00061-2
- Griffin, A. R., Potts, B. M., Vaillancourt, R. E., & Bell, J. C. (2019). Life cycle expression of inbreeding depression in *Eucalyptus regnans* and inter-generational stability of its mixed mating system. *Annals* of Botany, 124, 179–187. https://doi.org/10.1093/aob/mcz059
- Hajek, P., Kurjak, D., Von Wühlisch, G., Delzon, S., & Schuldt, B. (2016). Intraspecific variation in wood anatomical, hydraulic, and foliar traits in ten European beech provenances differing in growth yield. *Frontiers* in Plant Science, 7, 1–14. https://doi.org/10.3389/fpls.2016.00791
- Hardner, C. M., & Potts, B. M. (1995). Inbreeding depression and changes in variation after selfing in *Eucalyptus globulus* ssp. globulus. *Silvae Genetica*, 44, 46–54.
- Hedde, M., Aubert, M., Bureau, F., Margerie, P., & Decaëns, T. (2007). Soil detritivore macro-invertebrate assemblages throughout a managed beech rotation. Annals of Forest Science, 64, 219–228. https://doi. org/10.1051/forest
- Hedrick, P. W., Hellsten, U., & Grattapaglia, D. (2016). Examining the cause of high inbreeding depression: Analysis of whole-genome sequence data in 28 selfed progeny of *Eucalyptus grandis*. The New Phytologist, 209, 600–611. https://doi.org/10.1111/nph.13639
- Heineman, K. D., Turner, B. L., & Dalling, J. W. (2016). Variation in wood nutrients along a tropical soil fertility gradient. *The New Phytologist*, 211, 440–454. https://doi.org/10.1111/nph.13904

- Herlihy, C. R., & Eckert, C. G. (2002). Genetic cost of reproductive assurance in a self-fertilizing plant. *Nature*, 416, 320–323. https://doi. org/10.1038/416320a
- Hills, R. (2020). *Pericopsis elata*. The IUCN Red List of Threatened Species 2020.
- Hodgson, J. G., Montserrat-Martí, G., Charles, M., Jones, G., Wilson, P., Shipley, B., Sharafi, M., Cerabolini, B. E. L., Cornelissen, J. H. C., Band, S. R., Bogard, A., Castro-Díez, P., Guerrero-Campo, J., Palmer, C., Pérez-Rontomé, M. C., Carter, G., Hynd, A., Romo-Díez, A., De Torres Espuny, L., & Royo Pla, F. (2011). Is leaf dry matter content a better predictor of soil fertility than specific leaf area? *Annals of Botany*, 108, 1337–1345. https://doi.org/10.1093/aob/mcr225
- Hufford, K. M., & Hamrick, J. L. (2003). Viability selection at three early life stages of the tropical tree, *Platypodium elegans* (Fabaceae, Papilionoideae). *Evolution*, 57, 518–526. https://doi.org/10.1111/j. 0014-3820.2003.tb01543.x
- Ilunga-Mulala, C., Hatakiwe, H., Beeckman, H., Hardy, O. J., Ligot, G., Assumani, D., Ndjele, L., & Bourland, N. (2021). Influence of spacing and seed trees on the growth of *Pericopsis elata* saplings during the first twenty months of a planting trial. *Biotechnology*, *Agronomy*, *Society and Environment*, 25, 32–44.
- Ismail, S. A., Ghazoul, J., Ravikanth, G., Kushalappa, C. G., Uma Shaanker, R., & Kettle, C. J. (2014). Forest trees in human modified landscapes: Ecological and genetic drivers of recruitment failure in *Dysoxylum* malabaricum (Meliaceae). *PLoS One*, *9*, e89437. https://doi.org/10. 1371/journal.pone.0089437
- Jones, A., Breuning-Madsen, H., Brossard, M., Chapelle, J., Dampha, A., Deckers, J., Dewitte, O., Dondeyne, S., Gallali, T., Hallett, S., Jones, R., Kilasara, M., Le Roux, P., Micheli, E., Montanarella, L., Spaargaren, O., Thiombiano, L., Van Ranst, E., Yemefack, M., & Zougmoré, R. (Eds.). (2013). Atlas des sols d'Afrique. Commission européenne, Bureau des publications de l'Union européenne.
- Kafuti, C., Bourland, N., De Mil, T., Meeus, S., Rousseau, M., Toirambe, B., Bolaluembe, P. C., Ndjele, L., & Beeckman, H. (2020). Foliar and wood traits covary along a vertical gradient within the crown of long-lived light-demanding species of the Congo basin semideciduous forest. *Forests*, 11, 35. https://doi.org/10.3390/f1101 0035
- Kafuti, C., Van den Bulcke, J., Beeckman, H., Van Acker, J., Hubau, W., De Mil, T., Hatakiwe, H., Djiofack, B., Fayolle, A., Loubota Panzou, G. J., & Bourland, N. (2022). Height-diameter allometric equations of an emergent tree species from the Congo Basin. *Forest Ecology and Management*, 504, 119822. https://doi.org/10.1016/J.FORECO. 2021.119822
- Kouadio, Y., & Doucet, J. (2009). Etude du comportement de Baillonella toxisperma Pierre (moabi) dans les trouées d'abattage enrichies. Biotechnology, Agronomy, Society and Environment, 13, 317–324.
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). ImerTest package: Tests in linear mixed effects models. *Journal of Statistical Software*, 82, 1–26. https://doi.org/10.18637/JSS.V082.I13
- Laforest-Lapointe, I., Martínez-Vilalta, J., & Retana, J. (2014). Intraspecific variability in functional traits matters: Case study of Scots pine. *Oecologia*, 175, 1337–1348. https://doi.org/10.1007/s0044 2-014-2967-x
- Laure, N. M. R., William, M. A., & Din, N. (2014). Early growth stages structure and distribution of *Pericopsis elata* (Harms) van Meeuven in a logging concession of south-East Cameroon. *Journal of Biodiversity and Environmental Sciences*, 5, 354–363.
- Lavorel, S., & Garnier, E. (2002). Predicting changes in community composition and ecosystem functioning from plant traits: Revisiting the Holy Grail. Functional Ecology, 16, 545–556. https://doi.org/10. 1046/j.1365-2435.2002.00664.x
- Lenth, R. V., Banfai, B., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez, F., & Piaskowski, J. (2025). Package 'emmeans'. https://doi.org/10.1080/00031305.1980.10483031>. License

- Mosango, M. (1983). Une estimation du niveau d'accumulation des diaspores dans les sols des groupements herbacés à *Portulaca quadrifda* L. et à *Talinum triangulare* (Jacq.) Willd. à Kisangani (Zaïre). *Bulletin de la Societe Royale de Botanique de Belgique*, 116, 55–61.
- Murren, C. J., & Dudash, M. R. (2012). Variation in inbreeding depression and plasticity across native and non-native field environments. Annals of Botany, 109, 621–632. https://doi.org/10.1093/aob/ mcr325
- Naito, Y., Konuma, A., Iwata, H., Suyama, Y., Seiwa, K., Okuda, T., Lee, S. L., Muhammad, N., & Tsumura, Y. (2005). Selfing and inbreeding depression in seeds and seedlings of *Neobalanocarpus heimii* (Dipterocarpaceae). *Journal of Plant Research*, 118, 423–430. https://doi.org/10.1007/s10265-005-0245-z
- Nelder, J. A. (1962). New kinds of systematic designs for spacing experiments. Biometrics, 18, 283–307. https://doi.org/10.2307/2527473
- Ngongo, J. P. (2024). Inbreeding depression, functional traits and phenotypic plasticity in an endangered tree species from Congo basin with a mixed mating system [Data set]. Zenodo. https://doi.org/10. 5281/zenodo.13927593-01582-2
- Nickolas, H., Harrison, P. A., Tilyard, P., Vaillancourt, R. E., & Potts, B. M. (2019). Inbreeding depression and differential maladaptation shape the fitness trajectory of two co-occurring Eucalyptus species. Annals of Forest Science, 76, 10. https://doi.org/10.1007/s1359 5-018-0796-5
- Nicotra, A. B., Atkin, O. K., Bonser, S. P., Davidson, A. M., Finnegan, E. J., Mathesius, U., Poot, P., Purugganan, M. D., Richards, C. L., Valladares, F., & van Kleunen, M. (2010). Plant phenotypic plasticity in a changing climate. *Trends in Plant Science*, 15, 684–692. https:// doi.org/10.1016/j.tplants.2010.09.008
- Ouédraogo, D. Y., Fayolle, A., Daïnou, K., Demaret, C., Bourland, N., Lagoute, P., & Doucet, J. L. (2014). Enrichment of logging gaps with a high conservation value species (*Pericopsis elata*) in a central African moist forest. *Forests*, *5*, 3031–3047. https://doi.org/10. 3390/f5123031
- Oukabli, A., Lansari, A., Loudiyi, W., & Abousalim, A. (2001). Effets endogamiques sur la germination et la croissance de semis du cultivar autocompatible Tuono (*Prunus dulcis*). *Fruits*, *56*, 197–205. https:// doi.org/10.1051/fruits:2001123
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M. S., Cornwell, W. K., Craine, J. M., Gurvich, D. E., Urcelay, C., Veneklaas, E. J., Reich, P. B., Poorter, L., Wright, I. J., Ray, P., Enrico, L., Pausas, J. G., De Vos, A. C., ... Cornelissen, J. H. C. (2013). New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany*, *61*, 167–234. https://doi.org/10.1071/BT12225
- Pierce, S., Negreiros, D., Cerabolini, B. E. L., Kattge, J., Díaz, S., Kleyer, M., Shipley, B., Wright, S. J., Soudzilovskaia, N. A., Onipchenko, V. G., van Bodegom, P. M., Frenette-Dussault, C., Weiher, E., Pinho, B. X., Cornelissen, J. H. C., Grime, J. P., Thompson, K., Hunt, R., Wilson, P. J., ... Tampucci, D. (2017). A global method for calculating plant CSR ecological strategies applied across biomes worldwide. *Functional Ecology*, 31, 444–457. https://doi.org/10.1111/ 1365-2435.12722
- Pommerening, A., & Grabarnik, P. (2019). Individual-based methods in forest ecology and management. Springer International Publishing. https://doi.org/10.1007/978-3-030-24528-3
- Poorter, L., & Bongers, F. (2006). Leaf traits are good predictors of plant performance across 53 rain forest species. *Ecology*, 87, 1733–1743. https://doi.org/10.1890/0012-9658(2006)87[1733:LTAGPO]2.0. CO;2
- Poorter, L., Wright, S. J., Paz, H., Ackerly, D. D., Condit, R., Harms, E., Licona, J. C., Mazer, S. J., Webb, C. O., & Wright, I. J. (2008). Are functional traits good predictors of demographic rates? Evidence from five neotropical forests. *Ecology*, *89*, 1908–1920.
- Ratz, T., Perodaskalaki, A., Moorad, J., & Smiseth, P. T. (2020). Effects of inbreeding on behavioural plasticity of parent-offspring

interactions in a burying beetle. *Journal of Evolutionary Biology*, *33*, 1006–1016. https://doi.org/10.1111/jeb.13640

- Reed, D. H., Fox, C. W., Enders, L. S., & Kristensen, T. N. (2012). Inbreeding-stress interactions: Evolutionary and conservation consequences. Annals of the New York Academy of Sciences, 1256, 33–48. https://doi.org/10.1111/j.1749-6632.2012.06548.x
- Ren, L., Guo, X., Liu, S., Yu, T., Guo, W., Wang, R., Ye, S., Lambertini, C., Brix, H., & Eller, F. (2020). Intraspecific variation in *Phragmites australis*: Clinal adaption of functional traits and phenotypic plasticity vary with latitude of origin. *Journal of Ecology*, 108, 2531–2543. https://doi.org/10.1111/1365-2745.13401
- Richardson, A. E., & Simpson, R. J. (2011). Soil microorganisms mediating phosphorus availability. *Plant Physiology*, 156, 989–996. https://doi. org/10.1104/pp.111.175448
- Richardson, S. J., Allen, R. B., Buxton, R. P., Easdale, T. A., Hurst, J. M., Morse, C. W., Smissen, R. D., & Peltzer, D. A. (2013). Intraspecific relationships among wood density, leaf structural traits and environment in four co-occurring species of Nothofagus in New Zealand. *PLoS One*, 8, 24–26. https://doi.org/10.1371/journal.pone. 0058878
- Rozendaal, D. M. A., Phillips, O. L., Lewis, S. L., Affum-Baffoe, K., Alvarez-Davila, E., Andrade, A., Aragão, L. E. O. C., Araujo-Murakami, A., Baker, T. R., Bánki, O., Brienen, R. J. W., Camargo, J. L. C., Comiskey, J. A., Djuikouo Kamdem, M. N., Fauset, S., Feldpausch, T. R., Killeen, T. J., Laurance, W. F., Laurance, S. G. W., ... Vanderwel, M. C. (2020). Competition influences tree growth, but not mortality, across environmental gradients in Amazonia and tropical Africa. *Ecology*, 101, 1–11. https://doi.org/10.1002/ecy.3052
- Rozendaal, D. M. A., Hurtado, V. H., & Poorter, L. (2006). Plasticity in leaf traits of 38 tropical tree species in response to light; relationships with light demand and adult stature. *Functional Ecology*, 20, 207–216. https://doi.org/10.1111/j.1365-2435.2006.01105.x
- Sandner, T. M., Dotzert, A., Gerken, F., & Matthies, D. (2022). Inbreeding depression changes with stress response over time in flooded *Mimulus guttatus. Perspectives in Plant Ecology, Evolution and Systematics*, 57, 125697. https://doi.org/10.1016/j.ppees.2022. 125697
- Santiago, L. S., Wright, S. J., Harms, K. E., Yavitt, J. B., Korine, C., Garcia, M. N., & Turner, B. L. (2012). Tropical tree seedling growth responses to nitrogen, phosphorus and potassium addition. *Journal of Ecology*, 100, 309–316. https://doi.org/10.1111/j.1365-2745.2011. 01904.x
- Schlichting, C. D., & Levin, D. A. (1986). Effects of inbreeding on phenotypic plasticity in cultivated Phlox. *Theoretical and Applied Genetics*, 72, 114–119. https://doi.org/10.1007/BF00261465
- Schneider, C., Wayne, R., & Kevin, E. (2012). NIH image to ImageJ: 25 years of image analysis. Fundamentals of Digital Imaging in Medicine, 9, 671–675. https://doi.org/10.1007/978-1-84882-087-6_9
- Seltmann, P., Cocucci, A., Renison, D., Cierjacks, A., & Hensen, I. (2009). Mating system, outcrossing distance effects and pollen availability in the wind-pollinated treeline species *Polylepis australis* BITT. (Rosaceae). *Basic and Applied Ecology*, 10, 52–60. https://doi.org/10. 1016/j.baae.2007.11.008
- Silla, F., González-Gil, A., González-Molina, M. E., Mediavilla, S., & Escudero, A. (2010). Estimation of chlorophyll in *Quercus* leaves using a portable chlorophyll meter: Effects of species and leaf age. *Annals of Forest Science*, 67, 108. https://doi.org/10.1051/forest/ 2009093
- Smart, S. M., Glanville, H. C., Blanes, M. d. C., Mercado, L. M., Emmett, B. A., Jones, D. L., Cosby, B. J., Marrs, R. H., Butler, A., Marshall, M. R., Reinsch, S., Herrero-Jáuregui, C., & Hodgson, J. G. (2017). Leaf dry matter content is better at predicting above-ground net primary production than specific leaf area. *Functional Ecology*, *31*, 1336-1344. https://doi.org/10.1111/1365-2435.12832

- Stacy, E. A. (2001). Cross-fertility in two tropical tree species: Evidence of inbreeding depression within populations and genetic divergence among populations. *American Journal of Botany*, 88, 1041– 1051. https://doi.org/10.2307/2657086
- Sultan, S. E. (2000). Phenotypic plasticity for plant development, function and life history. *Trends in Plant Science*, 5, 537–542. https://doi. org/10.1016/S1360-1385(00)01797-0
- Swillen, I., Vanoverbeke, J., & De Meester, L. (2015). Inbreeding and adaptive plasticity: An experimental analysis on predator-induced responses in the water flea daphnia. *Ecology and Evolution*, 5, 2712– 2721. https://doi.org/10.1002/ece3.1545
- Takeuchi, Y., Kikuchi, S., & Diway, B. (2020). Albinism and inbreeding depression in seedlings of the tropical tree, *Shorea laxa. Journal of Forest Research*, 25, 413–419. https://doi.org/10.1080/13416979. 2020.1796897
- Taylor, S. H., Franks, P. J., Hulme, S. P., Spriggs, E., Christin, P. A., Edwards, E. J., Woodward, F. I., & Osborne, C. P. (2012). Photosynthetic pathway and ecological adaptation explain stomatal trait diversity amongst grasses. *New Phytologist*, 2, 387–396.
- Tyukavina, A., Hansen, M. C., Potapov, P., Parker, D., Okpa, C., Stehman, S. V., Kommareddy, I., & Turubanova, S. (2018). Congo Basin forest loss dominated by increasing smallholder clearing. *Science Advances*, 4, 1–12. https://doi.org/10.1126/sciadv.aat2993
- Valladares, F., Sanchez-Gomez, D., & Zavala, M. A. (2006). Quantitative estimation of phenotypic plasticity: Bridging the gap between the evolutionary concept and its ecological applications. *Journal* of Ecology, 94, 1103–1116. https://doi.org/10.1111/j.1365-2745. 2006.01176.x
- Velikova, V., Arena, C., Izzo, L. G., Tsonev, T., Koleva, D., Tattini, M., Roeva, O., De Maio, A., & Loreto, F. (2020). Functional and structural leaf plasticity determine photosynthetic performances during drought stress and recovery in two *Platanus orientalis* populations from contrasting habitats. *International Journal of Molecular Sciences*, 21, 1–18. https://doi.org/10.3390/ijms21113912
- Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., Jung, V., & Messier, J. (2012). The return of the variance: Intraspecific variability in community ecology. *Trends in Ecology* & *Evolution*, 27, 244–252. https://doi.org/10.1016/j.tree.2011.11. 014
- Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the concept of trait be functional! Oikos, 116, 882–892. https://doi.org/10.1111/j.2007.0030-1299.15559.x
- Vitasse, Y., Bresson, C. C., Kremer, A., Michalet, R., & Delzon, S. (2010). Quantifying phenological plasticity to temperature in two temperate tree species. *Functional Ecology*, 24, 1211–1218. https://doi.org/ 10.1111/j.1365-2435.2010.01748.x
- Walker, A. P., Beckerman, A. P., Gu, L., Kattge, J., Cernusak, L. A., Domingues, T. F., Scales, J. C., Wohlfahrt, G., Wullschleger, S. D., & Woodward, F. I. (2014). The relationship of leaf photosynthetic traits–Vcmax and Jmax–to leaf nitrogen, leaf phosphorus, and specific leaf area: A meta-analysis and modeling study. *Ecology and Evolution*, 4, 3218–3235. https://doi.org/10.1002/ece3.1173
- Wang, S., Li, Y., Ju, W., Chen, B., Chen, J., Croft, H., Mickler, R. A., & Yang, F. (2020). Estimation of leaf photosynthetic capacity from leaf chlorophyll content and leaf age in a subtropical evergreen coniferous plantation. Journal of Geophysical Research: Biogeosciences, 125, 1–14. https://doi.org/10.1029/2019JG005020
- Wang, Z., Townsend, P. A., & Kruger, E. L. (2022). Leaf spectroscopy reveals divergent inter- and intra-species foliar trait covariation and trait-environment relationships across NEON domains. New Phytologist, 235, 923-938. https://doi.org/10.1111/nph.18204
- White, F. (1986). La Végétation de L'afrique. ORSTOM-UNESCO.
- Winn, A. A., Elle, E., Kalisz, S., Cheptou, P., Eckert, C. G., Goodwillie, C., Johnston, M. O., Moeller, D. A., Ree, R. H., Sargent, R. D., & Vallejo-mar, M. (2011). Mixed-mating plants provides evidence for

16

selective interference and stable mixed mating. Evolution, 65(12), 3339–3359. https://doi.org/10.5061/dryad.gr43t

- Xiong, D., Chen, J., Yu, T., Gao, W., Ling, X., Li, Y., Peng, S., & Huang, J. (2015). SPAD-based leaf nitrogen estimation is impacted by environmental factors and crop leaf characteristics. *Scientific Reports*, 5, 1–12. https://doi.org/10.1038/srep13389
- Zhang, Q., Zhou, Z., Zhao, W., Huang, G., Liu, G., Li, X., & Wu, J. (2023). Effect of slope position on leaf and fine root C, N and P stoichiometry and rhizosphere soil properties in *Tectona grandis* plantations. *Journal of Forestry Research*, 34, 1997–2009. https://doi.org/10. 1007/s11676-022-01582-2

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Table S1. Correlation matrix between leaf and chemical traits, separately for outcrossed (top value printed in black) and selfed (bottom value printed in grey) seedlings.

Figure S1. Evolution of diameter (A–E) and total height (F–J) of Pericopsis elata plants for 4 years since transplantation across five competition levels (high to low from A–E and from F–J) for outbred (violet) and inbred (green) individuals.

Figure S2. Distribution of the values of nine leaf traits of *Pericopsis elata* during the first 4 years in plantation.

Figure S3. Variance partitioning of nine foliar traits (LFM, leaf fresh mass; LDM, leaf dry mass; LDMC, leaf dry matter content; LA, leaf area; SLA, specific leaf area; CHL, chlorophyll content; SS, stomatal size; SD, stomatal density; and Gmax, maximum stomatal conductance) across four explanatory variables for outbred (A) and inbred (B) trees.

How to cite this article: Ngongo, J. P., Kafuti, C., Beeckman, H., Ndjele, L., Ibrahim, E., Ilunga-Mulala, C., Hatakiwe, H., Brostaux, Y., Angbonda, D.-M., Drouet, T., Bourland, N., & Hardy, O. J. (2025). Inbreeding depression, functional traits and phenotypic plasticity in an endangered tree species with a mixed mating system. *Journal of Ecology*, 00, 1–17. <u>https://</u> doi.org/10.1111/1365-2745.70082