

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/352183981>

Human impact on symbioses between aquatic organisms and microbes

Article in *Aquatic Microbial Ecology* · August 2021

DOI: 10.3354/ame01973

CITATIONS

0

READS

104

16 authors, including:



Willem Stock

Ghent University

23 PUBLICATIONS 149 CITATIONS

[SEE PROFILE](#)



Martijn Callens

Centre d'Ecologie Fonctionnelle et Evolutive

15 PUBLICATIONS 304 CITATIONS

[SEE PROFILE](#)



Shira Houwenhuysse

KU Leuven

8 PUBLICATIONS 34 CITATIONS

[SEE PROFILE](#)



Ruben Schols

Royal Museum for Central Africa

8 PUBLICATIONS 35 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



PhD project The mud shrimp *Corophium volutator*: a key species in tidal flat sedimentary processes? [View project](#)



The influence of snail host microbiome in trematode parasite resistance [View project](#)



REVIEW

Human impact on symbioses between aquatic organisms and microbes

Willem Stock^{1,2,#}, Martijn Callens^{3,#}, Shira Houwenhuys⁴, Ruben Schols^{4,5},
Naina Goel^{4,6}, Manon Coone⁴, Charlotte Theys⁷, Vienna Delnat⁷, Alice Boudry^{4,7},
Ester M. Eckert⁸, Cecilia Laspoumaderes^{9,10}, Hans-Peter Grossart^{11,12},
Luc De Meester^{7,13}, Robby Stoks⁷, Koen Sabbe¹, Ellen Decaestecker^{4,*}

¹Laboratory of Protistology & Aquatic Ecology, Department of Biology, Ghent University, 9000 Ghent, Belgium

²Phycology Research Group, Department of Biology, Ghent University, 9000 Ghent, Belgium

³CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, 34080 Montpellier, France

⁴Laboratory of Aquatic Biology, KU Leuven Kulak, 8500 Kortrijk, Belgium

⁵Department of Biology, Royal Museum for Central Africa, 3080 Tervuren, Belgium

⁶Faculty of Bioscience Engineering, Department of Animal Sciences and Aquatic Ecology, University of Ghent, 8400 Oostende, Belgium

⁷Laboratory of Aquatic Ecology, Evolution and Conservation, KU Leuven, 3000 Leuven, Belgium

⁸Water Research Institute — National Research Council of Italy (CNR-IRSA), Molecular Ecology Group (MEG), 28922 Verbania, Italy

⁹Departamento de Ecología, Laboratorio de Limnología, INIBIOMA, CONICET-Universidad Nacional del Comahue, 8400 Bariloche, Argentina

¹⁰Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Biologische Anstalt Helgoland, 27498 Helgoland, Germany

¹¹Dept. of Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), 16775 Stechlin, Germany

¹²Dept. of Biochemistry and Biology, Potsdam University, 14469 Potsdam, Germany

¹³Dept. of Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), 12587 Berlin, Germany

ABSTRACT: Aquatic organisms rely on microbial symbionts for coping with various challenges they encounter during stress and for defending themselves against predators, pathogens and parasites. Microbial symbionts are also often indispensable for the host's development or life cycle completion. Many aquatic ecosystems are currently under pressure due to diverse human activities that have a profound impact on ecosystem functioning. These human activities are also expected to alter interactions between aquatic hosts and their associated microbes. This can directly impact the host's health and — given the importance and widespread occurrence of microbial symbiosis in aquatic systems — the ecosystem at large. In this review, we provide an overview of the importance of microbial symbionts for aquatic organisms, and we consider how the beneficial services provided by microbial symbionts can be affected by human activities. The scarcity of available studies that assess the functional consequences of human impacts on aquatic microbial symbioses shows that our knowledge on this topic is currently limited, making it difficult to draw general conclusions and predict future changes in microbial symbiont–host relationships in a changing world. To address this important knowledge gap, we provide an overview of approaches that can be used to assess the impact of human disturbances on the functioning of aquatic microbial symbioses.

KEY WORDS: Host–symbiont interactions · Aquatic microbial symbioses · Mutualism · Anthropogenic disturbances

*Corresponding author: ellen.decaestecker@kuleuven.be

#These authors contributed equally

1. INTRODUCTION

A large fraction of ecological interactions within an ecosystem are symbiotic, which can be defined as a phenomenon in which dissimilar organisms live together (De Bary 1879). Although the term symbiosis is mostly used in the context of mutualistic or commensalistic interactions between 2 species that live in close contact for a substantial amount of time, it encompasses a whole spectrum of outcomes, including parasitism. Furthermore, the outcome of a symbiosis between 2 species is not fixed, and both the strength and direction (positive, neutral or negative) of the interaction can change over time and/or shift depending on the context (Daskin & Alford 2012). Symbiosis can either be obligate, where one or both partners cannot survive without the other, or it can be facultative, with both partners able to survive independently outside of the symbiosis.

Microbial symbiosis refers to the phenomenon where, in general, a larger organism (host) is colonized by smaller unicellular microorganisms (symbionts). In aquatic environments, a wide diversity of eukaryote, bacterial and archaeal phyla engage in microbial symbiosis, either as host or symbiont (Grossart et al. 2013). Microbial symbionts can live as endosymbionts within the cells or as specialized organs of the host. However, many microbial symbionts colonize the surface of epithelial tissues such as the gut mucosa or are externally associated with the host, such as those growing in the phycosphere of microalgae (Zoccarato & Grossart 2019). The assembly of microorganisms found on a host is referred to as the host-associated microbiota, and they form—in combination with their specific habitat and biological activity—the microbiome (sensu Berg et al. 2020;

Fig. 1). This commonly used microbiome definition, however, is not fixed and often varies depending on the referenced source. There has been increased research interest in the host-associated microbiota, facilitated through new DNA sequencing technologies, which has revealed that many host organisms support highly diverse communities of microbial symbionts (Ley et al. 2008, White et al. 2016). In this review, we discuss both specialized endosymbionts and the broader host-associated microbiota, as we expect some parallels in how they interact with the host and how they are affected by environmental change. We only focus on microbial symbionts that provide—at least under some circumstances—a benefit to the host.

Most aquatic organisms start interacting with microbial symbionts from their birth, and sometimes even before (Bates et al. 2006, Nyholm 2020). Furthermore, aquatic organisms acquire bacteria from the environment throughout their life cycle, making the microbiota develop concomitantly with the host's chronological development. For several host organisms, it has also been shown that they exert control over the presence or abundance of specific symbionts, often favouring beneficial symbiont strains (Rawls et al. 2006, McFall-Ngai 2014, Tasiemski et al. 2015, Stock et al. 2019a). These processes often ensure that the microbial community provides the necessary functions throughout the host's life cycle (Sampson & Mazmanian 2015, Dominguez-Bello et al. 2019). Microbiota can be acquired from conspecifics through maternal transmission or other forms of interactions, or they can be acquired through colonization from the available pool of environmental microorganisms (Funkhouser & Bordenstein 2013). The mode of transmission is a key element in eco-evolutionary host–symbiont dynamics and is correlated with symbiont function and degree of specialization (Macke et al. 2017b). Many (especially aquatic) organisms obtain their microbiota through horizontal transmission, and the degree of exposure to environmental sources of microorganisms often plays a major role in determining microbiota assemblies (Adair & Douglas 2017). Intermediate modes, whereby symbionts from the parents are horizontally transmitted to their offspring, have also been observed (Ebert 2013, Björk et al. 2019). Vertically transmitted symbionts, which are transferred with high fidelity from the mother to the offspring




Fig. 1. Examples of aquatic host–symbiont associations. The host (largest organism, grey) with its symbionts (unicellular eukaryotes, *Archaea* and *Bacteria*, coloured). We define the symbionts as the microorganisms living in the host or externally associated with it. Well-studied examples of host–symbiont associations (orange) include (A) *Daphnia magna* and its gut microbiota, (B) *Euprymna scolopes* and the bioluminescent bacterium *Vibrio fischeri* and (C) *Acropora* spp. and the eukaryotic algae *Symbiodinium* spp.

from the parents are horizontally transmitted to their offspring, have also been observed (Ebert 2013, Björk et al. 2019). Vertically transmitted symbionts, which are transferred with high fidelity from the mother to the offspring

before birth, tend to resemble organelles, and their presence often has higher relevance for the host's fitness (Bright & Bulgheresi 2010, Fisher et al. 2017). Interestingly, aquatic host organisms seem to depend less on vertical transmission than terrestrial hosts (Russell 2019).

It has become clear that, for many organisms, the composition of their symbiont community has a major influence on various physiological processes and, ultimately, overall health. Variation in the gut microbiota composition is known to be caused by complex interactions between a range of factors such as environment, diet, age and medication (Falony et al. 2016, Callens et al. 2020). Alterations in microbiota community composition can have negative effects on the host and can lead to a so-called 'dysbiosis' linked to a variety of diseases, but the causality of this variation on host health is often poorly understood (Flandroy et al. 2018). In contrast, hosts can also be equally healthy when harbouring quite different microbiota, indicating that there is often a certain degree of functional redundancy between symbiont species or communities (Moya & Ferrer 2016, Callens et al. 2018).

Many aquatic ecosystems are currently under pressure due to diverse human activities such as land-use change, hydrological alterations to water bodies, pollution, deep-sea mining and commercial fishing. In addition, the effects of greenhouse gas emissions are expected to increasingly impact aquatic systems through climate change and ocean acidification (Doney et al. 2020). Although many research efforts are currently being undertaken to understand the potential impact of human activity on aquatic ecosystems and find ways to mitigate them, knowledge on the impact of human activity on microbial symbioses is scarce (Evariste et al. 2019, Timmis et al. 2019). This impact is, however, expected to have profound effects on aquatic ecosystems given the exceptional importance of microbial symbiosis for the health of many aquatic organisms (Cavicchioli et al. 2019). Effects of aquatic symbiotic interactions often extend far beyond the involved partners and can provide functions that are fundamental to whole aquatic ecosystems, such as energy provision to coral reefs (Muscatine 1990) and deep-sea hydrothermal vent communities (Jannasch & Mottl 1985), detoxification of seagrass bed sediments (van der Heide et al. 2012) or nutrient cycling in the water column, which can greatly affect primary production (Seymour et al. 2017).

The aim of this review is to assess the potential impact of human disturbances on interactions between aquatic organisms and their microbial symbionts, with a special focus on the consequences for the hosts'

health. In Section 2 ('Importance of the microbiome for aquatic organisms', summarized in Fig. 2A) we provide an overview of the importance of microbial symbionts for coping with various challenges encountered by aquatic organisms. We focus on resource acquisition, physiological stress, predation, pathogens, parasites and host development. In Section 3 ('Effects of anthropogenic disturbance on host-symbiont interactions', summarized in Fig. 2B and Table 1), we consider how the beneficial services provided by microbial symbionts can be affected by various types of human impacts on aquatic ecosystems: eutrophication, global warming, salinity changes, pollution and antibiotics. In Section 4 ('Perspectives'), we highlight the importance of including host-symbiont interactions when assessing the impact of human disturbances on aquatic ecosystems. We also identify some important knowledge gaps concerning the effect of human disturbances on the functioning of aquatic microbial symbioses and indicate how these knowledge gaps might be addressed in future studies.

2. IMPORTANCE OF THE MICROBIOME FOR AQUATIC ORGANISMS

2.1. Role of the microbiome in host development

Microbial symbionts have often been found to be essential for the development of multicellular aquatic hosts. Many hosts receive growth factors and vitamins required for their development from their microbiota (Sokolovskaya et al. 2020). Sea lettuce *Ulva mutabilis*, for example, requires multiple regulatory factors produced by its associated bacteria to develop into a 'blade' with rhizoids (Provasoli 1958, Wichard et al. 2015). In the absence of bacteria, sea lettuce develops into callus-like colonies consisting of undifferentiated cells. Several aquatic metazoan larvae (Porifera, Cnidaria, Acoelomorpha and Mollusca) require bacteria for their settlement (Tran & Hadfield 2011, Sneed et al. 2014, Fieth et al. 2016) whilst the presence of dinoflagellates belonging to the Symbiodinaceae seems to be required for successful metamorphosis in multiple marine organisms (Mies et al. 2017). In the sponge *Amphimedon queenslandica*, a bacterial symbiont belonging to the order *Chromatiales* supplies the amino acid L-arginine, which is essential for the sponge larvae to successfully settle and metamorphose (Fieth et al. 2016). Zebrafish *Danio rerio* display incomplete development and impaired function of their gastrointestinal tract in the absence of gut microbiota. This can, how-




Fig. 2. Schematic depiction of beneficial services provided by microbial symbionts and the potential impact of human activities on the interaction between aquatic hosts and their microbiota. (A) Important functions of the microbiota in pristine systems, including (from top, clockwise) defence against external toxins, provision of secondary metabolites, acquisition of resources, production of toxins, host development, predation defence, pathogen defence and osmoregulation. (B) Anthropogenic disturbances that can affect the host's microbiome, including (red arrows from top, clockwise) synthetic toxins, micro- and nanoplastic pollution, altered habitat structure (i.e. channelling of water bodies), increased abundance of antibiotic resistant and pathogenic microorganisms and increased temperature. The potential impacts of disturbing the microbiome on the host's health are indicated by grey arrows (from top, clockwise): reduced resource acquisition, increased pathogenic infections, lower toxin expulsion or degradation and reduced osmoregulation

ever, be reversed by inoculating them with their conventional microbiota, providing direct evidence for the role of the gut microbiota in gastrointestinal tract development (Bates et al. 2006). Furthermore, experimental evidence suggests that the microbiota is also required for a normal early life neurobehavioral development in zebrafish (Phelps et al. 2017).

Although less studied, associated microbiota can also play a role in the life cycle completion of unicellular aquatic hosts. This has been shown for the diatom *Seminavis robusta*, where its associated bacteria are known to affect sexual reproduction. They do this through modulation of the host's production of the sexual attraction pheromone diproline, which influences physical pairing of compatible cells and subsequent gametogenesis (Cirri et al. 2019).

2.2. Role of the microbiome in obtaining resources

Obtaining sufficient resources for growth, reproduction and meeting daily energy demands poses an

important challenge for all aquatic organisms. To cope with this challenge, many aquatic organisms rely on microbial symbionts that can either provide their host with organic carbon derived from primary production or with essential nutrients, enhance food digestion or help attract prey.

Autotrophic microbial symbionts often directly provide their aquatic host with energy derived from solar radiation (photosynthetic symbionts; Smith et al. 1969) or oxidation of electron donors (chemoautotrophic symbionts; Jannasch 1985). This relationship can result in a reduced dependency on external food resources for the host and may even result in an *in se* heterotrophic host obtaining all its required carbon from symbiotic primary producers (e.g. Hinzke et al. 2019). Symbiotic associations with photosynthetic microorganisms are widespread in the photic zone of aquatic environments. The endosymbiosis of the unicellular dinoflagellate *Symbiodinium* with cnidarian corals is a well-studied example of such an association (Freudenthal 1962). *Symbiodinium* provides oxygen and organic compounds to the coral and, in

Table 1. Potential functional impacts of anthropogenic disturbances on host–symbiont interactions

Anthropogenic disturbance	Functional impact of disturbance	Host	References
Changes in nutrient availability/eutrophication	Reduced calcification rates	Coral	van Oppen & Blackall (2019)
	Increased invasion of opportunistic pathogens	Coral	Voss & Richardson (2006), Gochfeld et al. (2012), Bourne et al. (2016)
Global warming	Dysbiosis of skin microbiota	Fish	Krotman et al. (2020)
	Increase in beneficial bacteria	Fish	Xie et al. (2011), Dong et al. (2013)
	Changes in the microbiome metabolism	Frog	Fontaine & Kohl (2020)
	Changes in digestive performance of the host	Salamander	Fontaine et al. (2018)
	Reduced exchange of symbiont-derived photosynthates	Coral	Baker et al. (2018)
Anthropogenic structures	Adaptation of symbionts to higher temperature and increased bioluminescence	Squid	Cohen et al. (2019)
	Increased susceptibility to disease	Bivalve	Li et al. (2019)
	Increase in epifauna and potential pathogens	Kelp	Marzinelli et al. (2009, 2018)
Salinity changes	Increased osmolyte production, sulphur oxidation and nitrogen fixation	Coral	Röthig et al. (2016)
	Reduced digestion of algal diet	Brine shrimp	Nougué et al. (2015)
	Changes in digestive and osmoregulatory capacity	Snail	Kivistik et al. (2020)
Chemical pollution	Increase in xenobiotic biodegradation capacity	Bivalve	Milan et al. (2018)
	Reduced body weight	Fish	Kan et al. (2015)
Antibiotics	Changes in ecological interactions within microbiota which affect host growth rates	Water flea	Callens et al. (2018)
	Reduced fitness under poor dietary conditions	Water flea	Akbar et al. (2020)
	Increased body weight and altered liver function	Fish	Keerthisinghe et al. (2020)
	Increased mortality when challenged with a pathogen	Fish	Zhou et al. (2018)
	Changes in host development	Fish	Yu et al. (2020)
	Impaired intestinal morphology, permeability and dysbiosis	Fish	Limbu et al. (2018)

turn, receives inorganic nutrients and protection from the host (Gordon & Leggat 2010). Other well-studied associations include the freshwater cnidarian *Hydra* with the green algae *Chlorella* (Ye et al. 2020), an association known since the 1920s (Goetsch 1924), and the sacoglossan sea slug *Elysia chlorotica* with the stramenopile algae *Vaucheria litorea* (West 1981, Chan et al. 2018). The latter association differs from the former since the host does not retain viable symbionts but merely the algal plastids. Many protists also harbour eukaryotic or prokaryotic microalgal symbionts that provide their host with photosynthates (Stoecker et al. 2009, Decelle et al. 2015, Leles et al. 2017). Furthermore, these photosynthetic symbionts often provide their host with additional biochemical functions (Nowack & Melkonian 2010). For instance, some foraminiferans host diatoms or dino-

flagellates, from which they obtain photosynthates, and simultaneously host cyanobacteria, which provide photosynthesis-dependent nitrogen (N) fixation (Lee 2006, Prazeres & Renema 2019).

Aquatic organisms that acquire energy from associated chemoautotrophic microorganisms are mostly found around deep-sea vents (Dubilier et al. 2008). Several tube-dwelling annelids nutritionally associate with aerobic methane-oxidizing bacteria (Goffredi et al. 2020), and the giant hydrothermal vent tubeworm *Riftia pachyptila* relies entirely on sulfide-oxidizing chemoautotrophic microorganisms for nutrition (Hinze et al. 2019). Symbiont-hosting vesicomyid clams can take up hydrogen sulfide from the sediment through their foot, and thioautotrophic bacteria that live in the gills of the clams can utilize this hydrogen sulfide as the main energy source (Lan et

al. 2019). Cold seep sponges from the genus *Suberites* receive energy from their archaeal symbionts, which carry out ammonia oxidation and carbon fixation within the sponge (Tian et al. 2017).

Gut microbiota play a vital role in food digestion for heterotrophic organisms. This holds true for many aquatic organisms, ranging from invertebrates such as crustaceans (Harris 1993, Callens et al. 2016, Holt et al. in press) to fishes (Egerton et al. 2018) and whales (Miller et al. 2020). For some organisms, bacteria provide essential enzymes to digest complex food sources. This is, for instance, the case in teredinid wood-boring bivalves — known as ship-worms — who depend on their associated bacteria to successfully digest wood (Sabbadin et al. 2018). Heterotrophic organisms also often rely on their gut microbiota to supply them with essential vitamins (Putnam & Goodman 2020).

Many aquatic phototrophs rely on their associated microorganisms to enhance the uptake of limited nutrients or to get access to nutrient resources that are otherwise unavailable to them. Through remineralisation and respiration, heterotrophic bacteria regenerate carbon dioxide out of leaked photosynthates, thereby facilitating nutrient circulation and increasing carbon availability for the host (Christie-Oleza et al. 2017). Algae-associated bacteria increase bio-availability of iron through siderophore production (Amin et al. 2009, Kurth et al. 2019), whilst N₂-fixing cyanobacteria, often as endosymbionts in microalgae, provide N in return for photosynthetically fixed carbon from the host (Villareal 1992, Foster & Zehr 2019). In addition to specific growth factors such as the auxin phytohormone indole-3-acetic acid (Seyedsayamdost et al. 2011, Amin et al. 2015), heterotrophic bacteria supplement various algae with vitamins (Croft et al. 2005, Sokolovskaya et al. 2020), complementing the host's nutritional needs.

An additional mechanism through which microbial symbionts can provide their heterotrophic host with food is by enhancing their capability to catch prey. This is the case in anglerfish, who host bioluminescent bacteria belonging to the genus *Photobacterium* in a light organ at the tip of their 'fishing rod' where they produce the necessary light to detect or attract prey (Hellinger et al. 2017, Michiels et al. 2018).

2.3. Role of the microbiome in dealing with physiological stress

Harsh environmental conditions can induce pronounced physiological stress in aquatic organisms.

Although the underlying mechanisms are not always evident, case studies suggest that associated microbiota can play a relevant role in reducing physiological stress in their host (White & Torres 2009).

Several unicellular aquatic hosts are known to gain protection against harmful solar radiation through their microbial symbionts. In the ciliate *Paramecium bursaria*, the presence of algal symbionts can reduce photo-oxidative stress caused by high UV radiation, most likely through the activity of antioxidants produced by the symbiotic algae countering reactive oxygen species production (Hörtnagl & Sommaruga 2007). In the marine benthic ciliate *Maristentor diniferus*, mycosporine-like amino acids produced by a symbiotic *Symbiodinium* are also known to minimize damage from exposure to solar UV radiation (Sommaruga et al. 2006). Furthermore, endosymbiotic *Symbiodinium* in corals are known to gain protection not only against light but also thermal stress from their associated bacteria that produce the carotenoid zeaxanthin, which potentially mitigates the effect of environmental stress due to its antioxidant activity (Motone et al. 2020).

Many heavy metals are essential micronutrients for aquatic organisms but can become highly toxic at higher concentrations (Morel & Price 2003). A plethora of host-associated bacteria have been found to carry heavy metal resistance factors on mobile genetic elements and could indirectly confer heavy metal resistance to the host (Selvin et al. 2009, Chaturvedi et al. 2015). Bacteria associated with the green alga *Enteromorpha compressa* demonstrated a high copper tolerance and were suggested to play a role in reducing the negative impact of copper on the algae (Riquelme et al. 1997). Similarly, increased copper tolerance was observed in the cnidarian *Hydra* when it was associated with microalgal symbionts (Karntanut & Pascoe 2005). However, the exact mechanisms by which the symbionts increase copper tolerance in their host were not evident. The mussel *Bathymodiulus* sp., living in metal-rich hydrothermal vent environments, harbours symbiotic bacteria that can absorb metal ions and subsequently excrete the metals in particulate forms, thereby detoxifying heavy metals for their host (Hardivillier et al. 2004). Kayath et al. (2019) showed that intestinal bacteria isolated from guppy fish living in hydrocarbon- and trace metal-contaminated wastewater could tolerate trace metals such as Hg, Co, Zn and Pb to a higher degree. Additionally, many of these bacteria were able to degrade gasoline or diesel fuel hydrocarbons. These studies indicate that the associated microbiota potentially have an important role in mitigating the

effects of various forms of environmental stresses such as pollution in many aquatic organisms.

The associated microbiota can also protect the host against physiological stress induced by secondary compounds present in the food. For example, in the water flea *Daphnia magna*, tolerance to the toxic cyanobacterium *Microcystis aeruginosa* is mediated by its gut microbiota (Macke et al. 2017a). Moreover, cyanobacterial tolerance was shown to be dependent on multiple microbial interactions within the *D. magna* host (Boudry et al. 2020). Given that toxin production is commonly found in various freshwater and marine microalgae that cause harmful algae blooms, symbiont-mediated protection against these toxins might be important for many aquatic hosts.

Microbial symbionts have also been shown to provide protection against osmotic stress in several hosts. In response to high salinity levels, the algal symbiont *Symbiodinium* produces high levels of the osmolyte 2-O-glycerol- α -D-galactopyranoside (floridoside), thereby increasing the capacity of the coral to cope with the effects of osmotic stress (Ochsenkühn et al. 2017). For the seaweed *Ectocarpus*, bacteria are essential to make the switch from marine to freshwater conditions. Without its bacteria, the algal host does not survive the transition to freshwater, indicating that bacteria can provide essential functions to mitigate the negative effects of salinity changes (Dittami et al. 2016).

2.4. Role of the microbiome in the host defence against natural enemies

Aquatic organisms have to cope with various natural enemies in their environment, including predators, grazers and parasites. As a consequence, many aquatic organisms have symbiotic interactions with microorganisms that protect them against these enemies. Defensive symbioses exhibit a variety of mechanisms through which the host gains protection against its natural enemies (Clay 2014).

Defensive symbioses against predators and grazers can be mediated through symbiont-derived secondary metabolites that render the host unpalatable or toxic. This mechanism is well known from sponges and tunicates, which rely heavily on chemical defences (Flórez et al. 2015). For example, the dictyoceratid sponge *Dysidea herbacea* is host to the intracellular symbiotic cyanobacterium *Oscillatoria spongelliae* that produces various halogenated compounds such as dysideathiazole, which are fish-feeding deterrents, protecting the sponge against predation (Ridley et al.

2005). Recently, it was shown that *Haliclona* sp. sponges have intracellular renieramycin-producing bacteria with a strongly reduced genome in specialized chemobacteriocytes, indicating that both partners evolved a highly specialized symbiosis (Tianero et al. 2019). These renieramycins are known to be highly cytotoxic and are hypothesized to protect the sponge against predators and pathogens. Tetrodotoxin (TTX) is a neurotoxin that is frequently used by many aquatic organisms to protect themselves against predation. Organisms containing TTX span a wide range of taxonomic groups such as red algae, pufferfish, blue-ringed octopuses, crabs, starfish, flatworms and amphibians (Jal & Khora 2015) and have evolved resistance through modification of their sodium channels (Venkatesh et al. 2005, Vaelli et al. 2020). Although this toxin can be accumulated by feeding on TTX-containing prey, several studies have shown the importance of symbiotic TTX-producing bacteria for host defence (e.g. Noguchi et al. 1986, Vaelli et al. 2020). Interestingly, this toxin is produced by a wide array of bacterial phyla (*Actinobacteria*, *Bacteroides*, *Firmicutes*, *Proteobacteria*) associated with various host organisms (Jal & Khora 2015), which could explain its widespread occurrence as a defensive mechanism. In addition to relying directly on microbial symbionts for protection, some animals depend on symbioses between their prey and its symbionts for chemical defence. This relationship was recently shown by Zan et al. (2019) for the sacoglossan sea slug *Elysia rufescens* and the algae *Bryopsis* sp., which both rely on the toxic lipopeptide kahalalide F for chemical defence. *Bryopsis* sp. obtains this toxin by housing symbiotic kahalalide-producing intracellular bacteria. *E. rufescens*, in turn, relies on this symbiosis to extract and sequester kahalalide F through feeding on *Bryopsis* sp. that contain kahalalide-producing symbionts. This example nicely shows that host-symbiont interactions can have consequences that extend beyond the involved partners in their influence on aquatic food webs and ecosystems.

Another mechanism for defensive symbiosis against predators is through counter-illumination with the purpose of host camouflage. Ventrally directed luminescence produced by symbiotic bacteria disrupts the shadow of the host that is cast by light coming from above, providing camouflage to avoid detection from below. This type of symbiosis is found in several species of squid and marine fish that host bacteria from the *Vibrionaceae* family (*Vibrio* sp. or *Photobacterium* sp.) in specialized light organs (Dunlap et al. 2007). These bacteria are acquired from the environment, and their hosts have evolved ways to select only ben-

eficial symbionts to populate the light organ (McFall-Ngai 2014). This interaction was shown to be structured through quorum sensing between the bacterial strains (Verma & Miyashiro 2013). For several species of squid and leognathid fish, it has been shown that the hosts can regulate the intensity, colour, angular distribution and patterns of ventral light emission in response to environmental light conditions by modifying the tissues surrounding the light organ (McFall-Ngai & Morin 1991).

A peculiar form of a microbial symbiosis that confers protection against predation can be found in ciliates of the genus *Euplotidium* that live in tide pools along rocky shores. A well-defined cortical band on the dorsal surface of the ciliate host is often colonized by bacterial episymbionts, referred to as epixenosomes, that belong to the genus *Verrucomicrobia* (Petroni et al. 2000). After attachment to the host, epixenosome cells differentiate themselves to form a sophisticated extrusive apparatus that can eject a ribbon in response to external signals. It has been experimentally shown that colonization by epixenosomes effectively protects *Euplotidium* from ingestion by the ciliate predator *Litonotus* sp. (Rosati et al. 1999).

The associated microbiota of aquatic organisms often play a crucial role in the host's resistance or tolerance to pathogen infection (Kimura & Tomaru 2014, Dheilly et al. 2015, Greenspan et al. 2019, Davoodi & Foley 2020, Schellenberg et al. 2020, Portet et al. 2021). For instance, Huot et al. (2020) reported a congruence of vector snails' phylogeny and their associated microbiome, combined with variable susceptibility between snail species and even populations of the same species, suggesting that the snail's microbiome might play a role in parasite resistance. Microbiota-induced pathogen resistance or tolerance has 3 underlying mechanisms: competition for resources and space, production of antimicrobial substances and stimulation of the host's immune response (Dheilly et al. 2015, Mallon et al. 2015, Knutie et al. 2017).

An example of colonization resistance through resource competition can be found in the associated microbiota of the coral *Acropora palmata*. In this host species, both the commensal microbiota and the white-pox-causing pathogen *Serratia marcescens* are known to employ glycosidases and N-acetyl-glucosaminidase to utilize components present in the coral mucus. Krediet et al. (2013) showed that several members of the *A. palmata* microbiota have the ability to inhibit the induction of these catabolic enzymes in *S. marcescens*. This inhibition gives the pathogen a competitive disadvantage

when growing in coral mucus, which results in reduced virulence.

Antimicrobials produced by the associated microbiota can also play a crucial role in the pathogen resistance of the host. In the marine sponge *Erylus discophorus*, a large and diverse fraction of the associated microbiota is known to produce antimicrobial compounds (Graça et al. 2013). However, sometimes the presence or absence of one specific bacterial strain producing antimicrobial substances can determine pathogen susceptibility or resistance. For example, the presence of the bacterium *Janthinobacterium lividum* in the cutaneous microbiota of the mountain yellow-legged frog *Rana muscosa* leads to a significantly reduced mortality as a consequence of infections by the lethal fungus *Batrachochytrium dendrobatidis*. This bacterial strain is capable of producing the anti-chytrid metabolite violacein, effectively reducing the chytrid pathogen burden on the mountain yellow-legged frog (Harris et al. 2009).

The associated microbiome can also stimulate the host's immune response (Knutie et al. 2017, Murdoch & Rawls 2019). Knutie et al. (2017) showed that early-life disturbance of the bacterial communities associated with Cuban tree frog *Osteopilus septentrionalis* tadpoles results in a significantly higher infection by the parasitic worm *Aplectana hamatospicula* in adulthood. In contrast, disturbance of the associated bacterial communities of adult frogs did not result in a higher parasitic burden. This study indicates that the microbiota likely plays an important role in priming the immune system of juvenile frogs, protecting the host from parasites later in life. In zebrafish larvae, the immune response is boosted by an increase in microbiota complexity, but can also be affected by specific bacterial strains. By comparing germ-free to conventional zebrafish larvae, it became apparent that many of the immunity-related genes are microbiota-regulated (Murdoch & Rawls 2019).

It is worth noting that the diversity of the microbiota is often found to be positively correlated with colonization resistance, although a specific mechanistic explanation is often lacking. On the one hand, such a pattern could be caused by competition, as communities with high species richness occupy more niches and hence are more resistant to invading pathogens by leaving less space and nutrients available for the pathogen to exploit (Mallon et al. 2015). On the other hand, it could be that key species providing resistance (e.g. through the production of antimicrobial substances) are often missing in less diverse communities. A positive correlation between microbiota diversity and colonization resistance was

observed in the European common frog *R. temporaria*, which showed increased resistance to the emerging *Ranavirus* with a more diverse skin microbiome (Harrison et al. 2019). When the microbiota diversity in zebrafish is reduced through exposure to the antibiotic olaquinox, they also become more susceptible to infection by the pathogen *Aeromonas hydrophila* (He et al. 2017).

3. EFFECTS OF ANTHROPOGENIC DISTURBANCE ON HOST-SYMBIONT INTERACTIONS

3.1. Eutrophication and shifts in oxygen availability and nutrient ratios

Altered nutrient availability is one of the greatest disturbances for aquatic ecosystems globally. Excessive inflow of nutrients, mainly phosphorus (P) and N from different anthropogenic reservoirs, such as wastewater and agriculture, is a key factor in the process of eutrophication in lakes and coastal areas. The enhanced primary production resulting from eutrophication, in turn, encourages microbial activity and the consumption of dissolved oxygen in bottom waters and benthic habitats. The consequent hypoxia of these habitats is extremely destructive for aquatic life (Diaz & Rosenberg 2008, Rousi et al. 2019). Microbial responses to hypoxia are intimately tied to the geochemistry, in both sediments and the water column. Mat-forming microbes are a macroscopic feature of benthic marine ecosystems subjected to severe hypoxia. These mats are formed largely of filamentous sulfide-oxidizing bacteria. Many protozoans and metazoan animals live in association with mats of mega- and macro-bacteria, including some eukaryotes with symbiotic bacteria. A hypothesis is that large mat-forming, sulfide-oxidizing bacteria detoxify sediment by removing sulfide, and thus facilitate metazoan habitation (Levin et al. 2009). Changing temporal micro-environments with respect to oxygen depletion at alternating locations has been shown to exist, e.g. in sponges, and provides suitable conditions for the activity of its anaerobic microbial symbionts, fuelling the holobiont's metabolism (Lavy et al. 2016).

Increasing inputs of anthropogenic N and CO₂ from the atmosphere are also perturbing ocean nutrient levels, generating a progressive shift towards P or N limitation (Bindoff et al. 2019). Imbalanced carbon:nutrient ratios are known to strongly affect animal assimilation efficiencies, nutrient excretion, growth and reproduction, whether nutrients are in

deficit or excess (Darchambeau et al. 2003, Laspoumaderes et al. 2015, Zoccarato & Grossart 2019). Although understudied relative to classic ecological systems, it is reasonable to hypothesize that the diversity of an organism's microbiome might, at least partially, be governed by nutrient availability. For example, patchiness in microbial communities in the oligotrophic North Atlantic Ocean relies on microbial associations with copepods as a source of nutrients and shows synchronous changes with variability in copepod nutritional content. As a consequence, bacterial communities associated with copepods in the oligotrophic ocean may not face nutrient limitation to the same extent as the surrounding free-living community (Shoemaker et al. 2020).

Changes in nutrient availability could also alter microbiome composition or functioning. Some organisms, such as sponges, have a highly stable microbiome when changing from oligotrophic to eutrophic conditions (Gochfeld et al. 2012, Simister et al. 2012, Luter et al. 2014). The great functional diversity and physiological plasticity of their microbiomes might contribute to their high ability to survive under variable environmental conditions (Baquiran & Conaco 2018). On the other hand, other aquatic organisms, such as the cyanobacterium *Trichodesmium* sp. (Frischkorn et al. 2017), larvae of the starfish *Acanthaster* (Carrier et al. 2018) or the benthic foraminiferan *Amphistegina lobifera* (Prazeres et al. 2017), among others, were reported to have a dynamic microbiome composition in response to changing nutrients. In all these cases, the plasticity of their microbiomes or the capacity to acquire different symbionts may underpin their success in changing systems and could represent an advantage in determining their resilience under changing environmental conditions.

Other groups are less tolerant of nutrient fluctuations. In corals, for example, an increase in N disrupts the finely tuned equilibrium in their microbiome, exacerbating the impact of increasing temperatures, slowing down calcification (van Oppen & Blackall 2019) and promoting the invasion of opportunistic pathogens (Voss & Richardson 2006, Gochfeld et al. 2012, Bourne et al. 2016). Zalewski et al. (2011) suggested that the *Daphnia* microbial gut flora competes with the host for P. Nutrient availability seems to change the host's microbiome communities and, in turn, the microbiota (including pathogens) play a role in how nutrients are used by the *Daphnia* host (Duffy et al. 2012, Aalto et al. 2015, Reyserhove et al. 2017). Krotman et al. (2020) showed how sporadic nutrient pollution events drove fish skin communities to dysbiosis through the alteration of the microbiome

bacterial groups reducing fish health. Macronutrients seem to also influence the composition and activity of intestinal microflora in fish. P and N supplementation increased the bacterial diversity of the gut microbiome, but in this case, with a trend to stimulate beneficial bacteria (Xie et al. 2011, Dong et al. 2013).

3.2. Global warming

The rise in global CO₂ concentration since 2000 is about 20 ppm decade⁻¹ (NOAA 2021). As a result, the Earth is heating up rapidly, yet is at a significant distance from thermal equilibrium due to the large flux of thermal energy currently entering the world's water bodies (Huntingford et al. 2020). The increasing water temperatures are decreasing oxygen solubility and increasing respiration rates and therefore amplifying the hypoxic conditions resulting from eutrophication.

Changes in environmental temperatures can influence the host's microbiome through direct effects of temperature on the associated microbiota community, or indirectly through temperature effects on the host phenotype, which in turn influences the microbiota (Kohl & Yahn 2016, Li et al. 2018, Fontaine & Kohl 2020). Shifts in the microbiota community composition can arise through changes in the external microbial pool to which hosts are exposed, and through host-mediated differences (Fan et al. 2013, Seedorf et al. 2014, Sullam et al. 2018). Indeed, differences in the effects of temperature on microbial community composition have been described under similar conditions between genotypes of the same species (Sullam et al. 2018, Frankel-Bricker et al. 2020) and between closely related species (Fontaine & Kohl 2020). Indirect changes of the associated microbiota can occur through modifications in active and passive selectivity for particular microbes in the host and are expected for several reasons (Nishiguchi 2000, Webster et al. 2008). Warming alters many aspects of ectotherm physiology, such as immune function (Maniero & Carey 1997) and gut transit time (van Marken Lichtenbelt 1992), which can influence microbial community structure (Hooper et al. 2012, Kashyap et al. 2013). Increased temperatures may also accelerate host metabolic rates (Kirk et al. 2018), leading to an increased host demand for carbon (carbohydrates) compared to N (amino acids) (Bestion et al. 2019). This was, for example, supported by enrichment in 2 of the 3 microbial carbohydrate metabolism pathways and a general decrease in microbial amino acid me-

tabolism pathways under increased temperatures in frogs (Fontaine & Kohl 2020). The changes in host metabolism and physiology can destabilize associations between hosts and microbial symbionts. For instance, in corals *Orbicella faveolata*, warming made the associated *Symbiodinium* parasitic, resulting in higher retention of its photosynthates (Baker et al. 2018).

In general, increasing water temperatures do not seem to affect microbial alpha diversity in a wide range of aquatic organisms (e.g. the water flea *Daphnia magna*: Sullam et al. 2018, Frankel-Bricker et al. 2020; *Lithobates* frogs: Kohl & Yahn 2016, Fontaine & Kohl 2020). Yet several exceptions have been documented where increased water temperatures reduced (e.g. Huyben et al. 2018, Li et al. 2018) or increased (e.g. Li et al. 2019) the alpha diversity, both of which may be associated with negative effects for the host. A reduction of gut-associated bacterial taxa may decrease the stability of the gut microbial communities (Fan et al. 2013) and reduce the host's resilience under stressful conditions (Lozupone et al. 2012). For example, increasing water temperatures caused the genus *Reyranella*, which plays a role in immune function (Peng et al. 2019), to be largely absent from the gut of the bullfrog *L. catesbeianus* (Fontaine & Kohl 2020). Furthermore, in the salamander *Plethodon cinereus*, the abundance of the genus *Janthinobacterium*, which protects amphibians against the fungal disease *Chytridiomycosis*, decreased under rising temperatures (Fontaine et al. 2018). An increased alpha diversity may also negatively impact the host when it reflects proliferation of opportunistic pathogens. For example, Li et al. (2019) indicated that the increase in *Vibrio* and *Arcobacter* under heat stress in the mussel *Mytilus galloprovincialis* increased its susceptibility to diseases, thereby contributing to increased mortality.

Elevated water temperatures typically alter the associated microbial community composition, as has been documented for many organisms including algae (e.g. Webster et al. 2011, Stratil et al. 2013), aquatic invertebrates (e.g. mussels: Li et al. 2018, Li et al. 2019; rotifers: Eckert et al. 2021; water fleas: Sullam et al. 2018, Frankel-Bricker et al. 2020) and vertebrates (e.g. fish: Huyben et al. 2018; frogs: Kohl & Yahn 2016, Fontaine & Kohl 2020). For example, the gut microbial communities of tadpoles of the invasive bullfrog *L. catesbeianus* change more rapidly and have more temperature-dependent bacterial functional pathways in response to higher temperatures than those of the non-invasive green frog *L. clamitans* (Fontaine & Kohl 2020). Such microbially mediated mechanisms may contribute to invasive

species being often more phenotypically plastic than native species (Davidson et al. 2011, Houwenhuysen et al. 2018).

Rapid evolutionary changes of the associated microbiota in response to increased temperatures might mitigate the negative effects of global warming experienced by the host. *Vibrio fischeri* evolved an increased ability to colonize the *Euprymna scolopes* squid host in response to higher temperatures resulting in an increase in bioluminescence (Cohen et al. 2019). Studies on *D. magna* showed that host clones from thermally different geographic regions harboured microbial communities that differed in structure, despite being reared under similar lab conditions for many years. However, the response in the gut microbiome to an increased temperature was similar for clones from different geographic regions, and the water flea population was still an important factor in explaining microbiome variation, indicating the potential for rapid evolution (Frankel-Bricker et al. 2020).

Next to the heating effect, the absorbance of enhanced CO₂ concentrations causes acidification of surface waters, which is particularly problematic for calcifying organisms such as foraminifera, corals and mussels (Orr et al. 2005). Corals growing close to a natural CO₂ seep hosted less symbiotic bacteria than the same corals growing further from the seep (Morrow et al. 2015). In addition, bacteria linked with diseases and stress seem to be more dominant in corals growing at a reduced pH (Meron et al. 2011). Negative feedback as a result of changes in the associated microbial community caused by acidification seems to worsen the effects of increased CO₂ concentrations on corals. Other species might benefit from the increased CO₂ levels. The same study (Morrow et al. 2015) that compared corals growing close to seeps to those growing at control sites also reported more photosynthetic microbes in sponges growing near the seeps, potentially providing the sponges with increased nutritional benefits.

3.3. Anthropogenic structures

Human interventions have changed residence time in lotic systems and created entirely new lentic systems, including reservoirs and canals. Additionally, many artificial structures such as windmill farms, oil rigs and aquaculture farms provide hard substrates for organisms to adhere to and shelter where previously there was none. Microorganisms can have a potential role in their host's adaptation to anthropo-

genically created habitats. They could do so by conferring resistance in the host to heavy metals from antifouling agents used to coat the structures (see Section 2.3) and extending the host's niche to better fit artificial local conditions. Overall, the impact of anthropogenic structures on the host-associated microbiota has hardly been explored.

Man-made structures, such as riverbank reinforcements and pier pilings, differ markedly from natural substrates by having different physico-chemical properties, orientation, shading etc. and generally strongly impact biological communities (Ferrario et al. 2016, Phillips & Prestie 2017, Reyne et al. 2021). Due to their effects on environmental conditions and biota, these structures can be expected to influence the host-associated microbiota and their interactions with the host (e.g. Jani & Briggs 2018). Marzinelli et al. (2009, 2018) showed that the kelp *Ecklonia radiata* growing on artificial pier-pilings was covered by more epifauna, and more of its epiphytic bacteria were associated with macroalgal diseases compared to kelp growing on neighbouring natural rocky reefs. Lower levels of light on the pier due to the shading of the pilings were put forward as one of the probable causes of the observed differences. Although the kelp populations did not differ in photosynthetic capacity, these results suggest a negative impact of the artificial structures on the host. Artificial substrates tend to provide suitable opportunities for novel, often invasive, organisms to settle (Mayer-Pinto et al. 2015, Ros et al. 2016). The same goes for newly introduced bacteria, which can potentially establish more easily on anthropogenic substrates where there may be a lower biological diversity (Amalfitano et al. 2015). Artificial substrates might therefore provide opportunities for novel associations between hosts and microorganisms.

3.4. Salinity changes

Salinization of freshwater environments is expected due to the global rate of mean sea-level rise at an average rate of 3.2 ± 0.4 mm yr⁻¹ since 1993 (Nicholls & Cazenave 2010). In combination with reduced rainfall, sea-level rise causes saltwater intrusion into coastal freshwater environments, converting them to brackish environments (Neubauer 2013). In addition, salt application for de-icing of roadways has been recognized as a major source of chloride ions to groundwater, streams, rivers and lakes (Dugan et al. 2017).

Changes in salinity are likely to result in differences in the abundance of key microorganisms (Herlemann

et al. 2011), shifts in microbial metabolism (Neubauer 2013) and nutrient cycling (Marton et al. 2012). The host's internal environment can shift as a consequence of salinity changes, which in turn may affect the microbiome structure. Taxonomy-based functional profiling of the host-associated microbiota in the osmoconforming coral *Fungia granulosa* after long-term increased salinity exposure resulted in a shift in the bacterial community towards increased osmolyte production, sulphur oxidation and N fixation (Röthig et al. 2016). Also in osmoregulators, changes can be expected, e.g. when salmon move from freshwater to saltwater, they drink continuously to counteract water loss to the hyperosmotic environment, thereby increasing the luminal alkalinity to that of the surrounding medium. Additionally, novel pathogens, microbes and dietary items are encountered in the saltwater environment, which suggests major changes and unique profiles in the intestinal microbiota following movement to saltwater (Dehler et al. 2017).

Depending on their salinity tolerance, strict freshwater bacteria may be extinguished while saline-tolerant bacteria may survive and marine bacteria could immigrate. Salinity shifts therefore also favour habitat generalists with a broad salinity tolerance (Székely & Langenheder 2014). *Artemia* survive at very high salinity levels but have a strongly reduced fitness at reduced salinity. Nogué et al. (2015) showed that this reduction in fitness is caused by the impact of salinity changes on the host's gut microbiota. The gut microbiota, necessary for algal digestion, was shown to grow better at high than at low salinity, and its functionality might thus be compromised at low salinity. Further evidence for the microbiota-mediated effect of salinity on host fitness was provided by the observation that axenic *Artemia* do not have reduced fitness at low salinity when feeding on easily digestible food. Kivistik et al. (2020) artificially increased or decreased the salinity level in water containing *Theodoxus fluviatilis* snails that originated from freshwater or mesohaline environments. They found that an increase in salinity in the water containing freshwater snails resulted in a strong change in the associated bacterial community, and typical marine bacteria became more pronounced in the digestive tract. However, the composition of the digestive tract microbiome of mesohaline snails did not alter after either an increase or decrease in salinity. Analysis of the functional profile of the digestive tract microbiomes showed that mesohaline snails have the ability to maintain the original bacterial community with high cellulolytic potential and the ability to produce osmolytes, while freshwater snails

were not able to compensate for the loss of functions by a shift in the bacterial community (Kivistik et al. 2020). This finding suggests that changes in salinity can result in compositional changes in the gut bacterial community, which in turn changes its functional profile and can have an effect on the host's fitness.

3.5. Chemical pollution

Anthropogenic activities such as food production and industrial activity cause the release of various chemicals into the environment that can have a considerable impact on the biotic properties of natural aquatic ecosystems. There is clear evidence that anthropogenic chemical pollutants adversely affect aquatic ecosystems through ecosystem destruction, habitat modification, water chemistry alteration and direct addition or removal of species (Malmqvist & Rundle 2002, Schallenberg & Armstrong 2004, Englert et al. 2013).

Chemical pollution with microplastics, silver nanoparticles, pesticides and heavy metals in aquatic environments can also disturb the associated microbiota of aquatic hosts living in those environments. For example, Milan et al. (2018) found that the hepatopancreas microbiota composition of the Manila clam *Ruditapes philippinarum* was influenced by the variation in overall chemical pollutant concentrations in its environment. In addition, they observed an over-representation of several pathways involved in xenobiotic biodegradation in the microbiota, suggesting a potential detoxifying action of the microbiome that can have consequences for the host's susceptibility to environmental chemicals. Alterations in the host-associated microbiota induced by chemical pollution can, in turn, have further effects on host physiology, growth and survival (Evariste et al. 2019).

Microplastics, defined as plastic debris smaller than 5 mm (Moore 2008), are a widespread environmental pollutant in both freshwater and marine environments (Welden & Cowie 2016). These microplastics can be ingested by many aquatic animals and have been found to negatively affect both phytoplankton and fauna in aquatic ecosystems (Wang et al. 2019). Two recent studies showed that exposure of zebrafish *Danio rerio* to polystyrene microplastics has significant effects on the phylum-level composition and diversity of its gut microbiota community (Qiao et al. 2019, Wan et al. 2019). Qiao et al. (2019) observed a decrease in *Proteobacteria*, while *Fusobacteria* increased as a result of polystyrene exposure. Wan et al. (2019) observed a decrease in *Gammaproteo-*

bacteria and *Bacteroidetes*, while *Firmicutes* significantly increased. Both studies indicated that these polystyrene-induced shifts in gut microbiota could be associated with the observed dysbiosis causing gut inflammation, metabolic disorders, oxidative stress or neurotoxicity. Silver nanoparticles decreased *Beta-proteobacteriales* (mainly *Curvibacter* and *Undibacterium*) in the microbiomes of the freshwater planarian *Schmidtea mediterranea* (Bijnens et al. 2021). These studies show that the presence of synthetic micro- and nanoparticles can have pronounced effects on host–microbiota interactions.

Pesticides commonly used in agriculture are found to be widely present in aquatic ecosystems due to runoff (Sánchez-Bayo et al. 2016). Kan et al. (2015) found that exposure of goldfish *Carassius auratus* to the pesticide pentachlorophenol (PCP) was associated with an increased abundance of *Bacteroidetes*, especially members from the *Bacteroides* genus, and a decreasing *Firmicutes/Bacteroidetes* ratio. The abundance of 4 other gut microbial taxa was furthermore negatively correlated with this increase in *Bacteroidetes*. These PCP-induced changes in gut microbiota community structure were thought to play a crucial role in the reduced body- and liver weight of these fishes when exposed to PCP. In contrast, exposure to pesticides does not necessarily always affect microbiota communities or symbiotic interactions. For example, Knutie et al. (2017) showed that exposure of the Cuban tree frog *Osteopilus septentrionalis* to environmental concentrations of the herbicide atrazine only had minimal effects on its associated microbiota, and no effects on the host's susceptibility to infection with the chytrid fungus *Batrachochytrium dendrobatidis*.

Elevated concentrations of heavy metals are often found in aquatic ecosystems due to runoff from mining, agriculture or industrial waste. Dahan et al. (2018) exposed larval zebrafish to environmental concentrations of arsenic (between 10 and 100 ppb). They found that even at the lowest concentration of arsenic there were significant changes in the gut microbiota community structure. Although the authors suggested that arsenic exposure induced a dysbiosis in the zebrafish microbiota, health consequences of this shift in microbiota for the host were not measured. Similarly, changes in gut microbiota community composition were found when zebrafish were exposed to lead (Xia et al. 2018). Here, the authors also suggested that the observed changes in gut microbiota composition might impact the host's metabolism, although direct effects of lead exposure on the host could not be excluded.

3.6. Antibiotics

Antibiotics are commonly used to treat or prevent bacterial infections in humans and animals, and traces can be found in aquatic systems as a result of the absence of (or incomplete) removal from wastewater (Giger et al. 2003). Concentrations of these antibiotics are in the range of ng or a few $\mu\text{g l}^{-1}$, and it is considered even lower for coastal sea waters (Gothwal & Shashidhar 2015). However, some studies report concentrations in the mg l^{-1} range for sulphonamides (Bilal et al. 2020, Bojarski et al. 2020). At higher concentrations, antibiotics often have toxic or growth-inhibiting effects, e.g. on *Daphnia* (Wollenberger et al. 2000, Gorokhova et al. 2015) and fish (Bojarski et al. 2020). Even at very low doses, antibiotics can show effects on microbiota composition and diversity in aquatic animals (Callens et al. 2018) as they have been shown to act as signal molecules (Sengupta et al. 2013). Such effects of individual antibiotics might be more pronounced considering that the exposure to such compounds is likely continuous and composed of a highly variable cocktail of various antibiotics (Evariste et al. 2019). The actual impact of this contamination on the microbiota of animals in nature is unknown since few studies have tackled the effect of this low-dose exposure, and many of them used concentrations that are in the upper range or above environmentally relevant ones. Here, we only discuss studies using $<50 \mu\text{g l}^{-1}$ of antibiotics and their related effects on microbiota. Yet all of the currently available literature used only 2 model systems—*Daphnia* or fish—which renders a generalization of these results difficult.

The few available data suggest that the impact of low-dose antibiotics on diversity and community composition as well as the resilience of the host-associated microbiota depend on both the antibiotic itself and on the host: a long-term exposure of juvenile zebrafish to relatively low concentrations of tetracycline ($1 \mu\text{g l}^{-1}$) showed a shift of the microbial community and indicated an increase in microbiota diversity. Bacterial taxa benefiting from the presence of the antibiotic included *Fusobacter*, *Firmicutes* and *Bacteroidetes* (Keerthisinghe et al. 2020). Consequently, an increase in fish body weight which altered the liver function was observed (Keerthisinghe et al. 2020). Exposure to a slightly higher concentration of Rifampicin ($25 \mu\text{g l}^{-1}$) for about 6 d resulted in a strong shift in the composition of both skin and gut microbiota of the western mosquitofish *Gambusia affinis* and a strong decrease of microbial diversity and culturability. Culturability of bacteria quickly increased during

antibiotic exposure, indicating higher growth of the selected resistant strains. However, neither skin nor gut microbiota composition or diversity were resilient to antibiotic disturbance and still showed highly altered microbiota composition after 1 wk without antibiotics. On the other hand, in a study where larval fathead minnows *Pimephales promelas* were exposed to low levels of Triclosan (100 ng l^{-1}), the antibiotic had an immediate effect on community composition and diversity, but the gut community recovered from this disturbance after 2 wk of antibiotic absence. The different studies used different antibiotics, animals and recovery times, thus more standardized experiments would be needed to draw generalized conclusions on the resilience of the animal-associated microbiota after antibiotic-induced disturbances.

Sometimes the effect of low-dose antibiotics is not visible at the level of the microbiota composition. When zebrafish were exposed for 2 mo to oxytetracycline at $10 \mu\text{g l}^{-1}$, the microbiota diversity was not impacted and the community composition was rather similar to the one of the no-antibiotic treatment. However, the microbiota composition of the surrounding water community shifted strongly in the presence of the antibiotic (Almeida et al. 2019). This suggests that compared to free-living communities host-associated communities might in some cases be more resistant to antibiotic exposure. In a different study on zebrafish, the same antibiotic at even lower concentration (420 ng l^{-1}) did shift the microbial community, but no shift was observed with sulfamethoxazole (260 ng l^{-1}) (Zhou et al. 2018).

Despite these small shifts in microbiota community composition, the presence of antibiotics can also affect the host's physiology or susceptibility, either through the direct action of the antibiotic on the host or indirectly through impacts on the microbiome. In the above-mentioned study, for example, the presence of both antibiotics was associated with higher mortality of the fish when challenged with a pathogen (Zhou et al. 2018). The same oxytetracycline, also in low dose, had a very strong effect on the development of the animals in other experiments (Yu et al. 2020), indicating a high potential of the antibiotic to affect the fish's physiology via changes in its microbiota. Exposure of Nile tilapia *Oreochromis niloticus* to low doses of sulfamethoxazole and oxytetracycline only slightly changed microbial community and diversity, but chronic exposure to antibiotics impaired intestinal morphology, permeability and induced microbiota dysbiosis with a very strong negative impact on the fish's physiology, nutritional metabolism and immune system (Limbu et al. 2018).

Moreover, combined effects might play an important role: in an experiment with the *D. magna* model system and low concentrations of ciprofloxacin, an effect on the microbiome composition (with increased abundances of *Pseudomonas*-related taxa) and on animal fitness was only observed when the diet was poor. No effect, however, was detected when animals were grown on a healthy diet (Akbar et al. 2020). Effects of antibiotics on the animal microbiota at very low concentrations ($<10 \text{ ng l}^{-1}$), found in many central European freshwater systems (Szymańska et al. 2019), are so far unstudied. Furthermore, sub-inhibitory concentrations of antibiotics often modulate bacterial gene expression, and many antibiotics are assumed to act as signalling molecules in microbial communities (Yim et al. 2007, Sengupta et al. 2013). As such, even in the absence of bactericidal or bacteriostatic effects, antibiotics might affect the physiology of microbial symbionts and ecological interactions between them. This, in turn, can affect the benefit they provide to their host. Thus, long-term exposure experiments with low antibiotic concentrations and additional model systems should be a focus of future research.

4. PERSPECTIVES

4.1. Significance of human-driven symbiotic shifts

The ever-increasing list of examples of beneficial effects of microbial symbionts on aquatic organisms indicates that our current knowledge on this topic is mainly limited by research efforts and that much remains to be discovered. Despite the overwhelming evidence for the importance of microbial symbiosis for aquatic organisms, we know very little about how human activities affect these symbioses, and what the consequences are for the hosts. Furthermore, studies investigating human impacts on microbial symbioses are often descriptive, reporting only changes in the presence or abundance of microbial symbionts without inferring fitness consequences for the host. The fact, however, that many descriptive studies often do report large shifts in the microbiota community suggests that disturbances in mutualistic interactions can be expected. This is supported by a number of studies — albeit limited — that found a significant impact of human activities on the symbiotic interaction itself (Table 1).

Obtaining a better insight into the role of the microbiome is crucial, given that it affects human and, more generally, ecosystem health as indicated

in the 'One Health' framework, where human and environmental health are considered to be interconnected and interdependent (Flandroy et al. 2018, Berg et al. 2020). Ecosystem health depends to a large degree on intact relationships between hosts and their microbiomes or symbionts. Major issues to be resolved in order to evaluate the importance of such symbioses for ecosystem health include evaluating how taxonomically/functionally redundant microbiota are with respect to their impact on host fitness (Koedooder et al. 2019) and what the ecosystem-level effects are, e.g. if keystone species (such as *Daphnia*) are impacted (Macke et al. 2020). In line with this 'One Health' framework, it has become apparent that integrating the interaction between multiple symbionts, their host and the environment is crucial for understanding disease aetiology and ecosystem health (Bass et al. 2019). Many infections have zoonotic reservoirs from which they can (re-)emerge (J. P. Webster et al. 2016, Leger & Webster 2017), while others may even pose conservation threats for wildlife populations (Thompson et al. 2010, Heard et al. 2013). Although the link between host health and its microbiota has mainly been considered in a medical context, its importance in other areas such as conservation biology is becoming increasingly clear (Trevelline et al. 2019). One way that humans have triggered wildlife declines is by transporting disease-causing agents to remote areas of the world (Byrne et al. 2019). Considering the importance of a host's microbiome in disease resistance and host health, one would expect the microbiome of aquatic hosts to play an important role in their resistance to and spread of infections (Toledo & Fried 2011). This role has been shown for the resistance of amphibians to the fungal pathogen *Batrachochytrium dendrobatidis* (Flandroy et al. 2018, Greenspan et al. 2019). The realization that mass drug treatments alone do not suffice to control diseases renewed the focus on the control of their vectors. Similar to malaria control, manipulating the microbiome of aquatic vectors could be a sustainable way of controlling these diseases without negatively affecting local ecosystems. As a first step, resistance-inducing strains could be identified, such as the *Enterobacter* bacterium in the mosquito *Anopheles gambiae* which

makes its host almost completely resistant to *Plasmodium* infections (Cirimotich et al. 2011). Such research is especially relevant in the Anthropocene as human activities alter ecosystems, reduce the microbial diversity in the environment and spread disease-transmitting species across the world (Houwenhuysse et al. 2018, Lachnit et al. 2019, Vanhove et al. 2020).

4.2. Future challenges

In this last section, we postulate 5 major challenges to be addressed in future microbiome–host studies in relation to anthropic pollution stresses (summarised in Fig. 3):

4.2.1. How can we distinguish direct effects of human activities on the host from indirect effects through their impact on symbiotic interactions?

One of the main challenges when assessing the effects of anthropogenic disturbances on symbiotic interactions is the difficulty in separating this impact from direct effects on the host (Fig. 3A). Indeed, direct effects on aquatic hosts have been recorded for all of the above-mentioned disturbances (e.g. Silvestre et




Fig. 3. Challenges to tackle in aquatic microbial research to improve our understanding of anthropogenic effects on symbioses between aquatic organisms and microbes. Over time, increasing anthropogenic pressures on an aquatic host (from left to right) inflict stress on the host and change its associated microbiota. The identified knowledge gaps are: (A) Do stressors directly impact the host or indirectly through the microbiota? (B) How do multiple stressors simultaneously impact the symbiotic interactions? (C) How do human activities impact microbial symbiont transmission? (D) What are the long-term effects of anthropogenic pressure on the symbioses and host persistence? (E) Can findings be generalised to other species?

al. 2012). Two methodological approaches can be used to separate such direct from indirect effects.

In a first approach, negative impacts on symbiotic interactions are indirectly inferred based on a profound knowledge of the functioning of the symbiosis, and on observations of changes in parameters that are known to be important for the symbiosis. This can be relatively straightforward if the benefit provided by a specific symbiont is well-described and the anthropogenic disturbance shows a clear effect on the presence, abundance or functioning of this symbiont (e.g. coral bleaching under temperature stress; Glynn 1991). Data on the effect size of symbionts on the host's phenotype through experiments manipulating their presence or abundance (e.g. Rosati et al. 1999, Becker et al. 2009) can furthermore be used to examine if the effect sizes of the human impact on the host are in the same range. However, indirect inference of negative effects on host–symbiont interactions is often not as straightforward because a benefit provided by symbionts can depend on complex interactions, and anthropogenic disturbances can impact microbiota communities in complex ways through changes in composition and metabolic functioning. The use of multi-omics approaches could provide insight into the magnitude of the impact of a disturbance on symbiotic interactions. For example, if the gut microbiota community benefits the host through the production of a specific metabolite; (meta)genomics, (meta)transcriptomics, (meta)proteomics and (meta)metabolomics could be used to investigate the impact of a disturbance on, respectively, the presence of genes needed for the production of this metabolite, the expression of these genes and the concentration of gene products or the metabolite itself (Douglas 2018, He et al. 2020).

The second approach involves directly measuring the effect of a disturbance in the associated microbiota on the host's phenotype. When an environmental factor causes a change in the presence or abundance of symbionts, this change can be mimicked in the host without exposing this host to the factor responsible for this change. This process allows the effect of changes in the microbiota community on the host phenotype to be measured by removing the direct effect of the environmental factor on the host. Two methods can be used to achieve this, both of them starting with an initially germ-free host, followed by the inoculation of microbial symbionts. In transplant experiments, hosts and their associated microbiota are exposed for some time to a stressor (or a combination of stressors) of interest (e.g. high temperature, toxic cyanobacteria, pesticides), allowing

the microbiota to respond to this change. Microbiota communities are subsequently extracted from the exposed hosts and transplanted into germ-free recipient hosts. This approach has been successfully used to directly measure the effect of environmentally induced changes in the microbiota on phenotypic traits of various animals such as mice, desert woodrats and *Daphnia* water fleas (Ridaura et al. 2013, Kohl et al. 2014, Macke et al. 2017a). Synthetic communities can also mimic the effect of a disturbance on the associated microbiota through precise inoculation of cultivated microbiota members. Although this method is still in its infancy due to a lack of representative collections of microbiota isolates, this approach is becoming feasible for some model organisms like *Caenorhabditis elegans* (Rafaluk-Mohr et al. 2018, Dirksen et al. 2020) and *Drosophila melanogaster* (Douglas 2018). Both methods hold great promise for directly measuring the effect on changes in the microbiota composition on the host phenotype, but it must be noted that accurately mimicking environmentally induced changes in the absence of the environmental factor can be challenging (Callens et al. 2018). Furthermore, metabolic changes in the microbiota as a result of removing the environmental factor of interest are also not taken into account.

4.2.2. Can we expect that the measured impact of single stressors will be additive in multi-stress environments?

The impact of humans on aquatic systems is clearly multifaceted, and aquatic organisms are therefore likely to face different challenges simultaneously (Fig. 3B). Examples have shown that stressors can amplify one another (e.g. acidification and temperature increase; N. Webster et al. 2016), but more complex effects are equally possible. For instance, a high salinity tolerance in a coral–*Symbiodinium* system also confers increased thermotolerance (Gegner et al. 2017), and the presence of cyanobacteria affects white fat cell disease in *Daphnia*, potentially mediated through changes in the microbiome (Coopman et al. 2014, Macke et al. 2017a). The complexity of multi-stressor interference on host–microbiome interactions calls for more realistic experiments evaluating the effects of multiple stressors on aquatic organisms simultaneously. Such a multi-stressor experimental approach will provide a more accurate insight into the performance of aquatic organisms in an anthropogenically disturbed environment.

4.2.3. What is the impact of human disturbances on microbiota transmission?

Very little is known about the consequences of an acute disturbance on symbiont transmission (Fig. 3C). Many of the examples discussed above have shown that anthropogenic disturbances can severely affect the presence or abundance of beneficial symbionts. These changes can subsequently disturb the symbiont transmission chain, affecting the transmission probability of beneficial symbionts to the next generation (Koskella & Bergelson 2020). This, in turn, can strongly affect subsequent assembly of host-associated microbiota communities and the host's phenotype (Callens et al. 2018). In laboratory populations of mice, a 'legacy effect' is often observed, where differences in microbiota composition are maintained between groups of co-housed individuals who transmit microbiota among each other but not with members of other groups (Robertson et al. 2019). This observation indicates that changes in the microbiota of an interacting population of hosts can be maintained over time, even if the stressor is no longer present.

Ecological frameworks such as metapopulation theory (e.g. used to understand the transmission of microbial infections; Gandon et al. 1996, Haag & Ebert 2004, Laine & Hanski 2006) and metacommunity theory (Leibold et al. 2004, Mihaljevic 2012) can be used to estimate the consequences of an anthropogenic impact on symbiont transmission between hosts, and the consequences of changes in transmission on host-associated symbiont communities. Microbiota studies can be perfectly treated as a metacommunity ecology problem, i.e. to adapt methods from metacommunity ecology to understand dispersal, diversity patterns and community assembly of symbionts (Macke et al. 2017b).

4.2.4. What is the potential for long-term mitigation of human impacts on symbiotic interactions?

Most environmental disturbances have marked and reasonably well-studied short-term effects on the interactions which are generally reflected by shifts in the host-associated microbial community as well as changes in the host's metabolism and physiology. The resulting changes in the symbiotic interactions can lead to dysbiosis (e.g. Krotman et al. 2020), worsening the impact of anthropogenic changes on the host, or shift the interaction towards a novel equilibrium between host, microbiota and the environ-

ment (e.g. Röthig et al. 2016). The long-term stability of this newly attained equilibrium has hardly been explored, yet is likely to determine how the hosts will endure the unremitting human pressure on aquatic ecosystems (Fig. 3D).

Understanding the ecological dynamics, whereby the symbiotic community changes as a result of anthropogenic impacts, and the evolutionary dynamic, a product of the adaptations of the host and its associated symbionts to anthropogenic impacts, is key to predicting long-term stability of the host-microbiota interactions and host persistence. Some host-microbiota systems might inherently be more robust to changes, either due to higher plasticity of the host, as is the case for some invasive species (e.g. Fontaine & Kohl 2020), or due to a more diverse associated microbial community with a high degree of functional redundancy (Callens et al. 2018). Anthropogenically induced environmental changes could, in these robust systems, result in a microbial community with a very similar functional make-up, providing similar health benefits for the host.

Host-microbiota systems with a less diverse microbial community or highly coevolved partners, on the other hand, are likely to be more sensitive to perturbations as replacement or complementation of the symbionts might not be possible. Rapid adaptations of the host, the microbiota or both might still aid these systems to persist under anthropogenic pressure.

If evolutionary rescue occurs, enabling the host and its associated microbiota to recover from anthropogenic pressure, it is likely driven by the evolutionary potential of the microbial community, which is generally much higher than that of the host (Koskella et al. 2017, De Meester et al. 2019). It is worth noting that adaptation of the microbial symbionts does not need to happen in association with the host but can even occur independently and still have implications for the health of the host. For instance, the adaptation of the bacterium *Vibrio fischeri* to temperature and pH stress affected its bioluminescence levels when present in its squid host (Cohen et al. 2019, 2020).

Long-term persistence of the organisms does not only depend on how changes in host-microbiota interactions will affect the health of the host, but also on how they will impact the fitness of the host compared to the other organisms present. Studies comparing how changes in the microbial symbiont community affect multiple competing species are rare for aquatic systems (Koedooder et al. 2019). Such studies can help explore how microbiome-driven changes in relative fitness can, in the long-term, result in alternate communities (terHorst & Zee 2016).

4.2.5. Can we generalize the anthropogenic impact on symbioses between aquatic organisms and symbionts?

Most studies are limited to a restricted set of organisms that either show drastic effects when microbial symbionts are disturbed (e.g. coral bleaching in the coral–zooxanthellae symbiosis; Muller-Parker et al. 2015) or freshwater organisms that are easily amenable to experimentation (e.g. *Hydra*, Deines & Bosch 2016 and *Daphnia*, Sison-Mangus et al. 2015, Callens et al. 2016, Macke et al. 2017b). Considering the large aquatic diversity which is, particularly at broad taxonomic scales, much higher than in terrestrial systems (Grosberg et al. 2012, Tadesse 2018), the limited set of focal organisms does not allow us to draw general conclusions upon how the disturbance of microbial symbioses might influence aquatic ecosystem functioning (Fig. 3E). This urges us to diversify the experimental systems in which these interactions are investigated. Broadening the scope of suitable host–microbiota systems poses important challenges, including developing the appropriate genetic tools and establishing axenization protocols, but major efforts are currently being made at overcoming these hurdles for a wide array of aquatic organisms (Dittami et al. 2021).

Generalization is further hampered by limited insight into how genetic diversity within species impacts the host–microbiota interactions. It is clear that different genotypes of the same species can have markedly different microbiomes (e.g. Callens et al. 2020, Frankel-Bricker et al. 2020), respond differently to environmental changes (Stock et al. 2019b) and affect the environmental microbial communities differently (Macke et al. 2020, Massol et al. 2021). Studies using a single genotype might thus not be representative of the response of genetically diverse species, hampering generalization of human impacts even on the level of a single species. A relevant discussion in this respect is whether there are consistent patterns to be detected between field and lab microbiomes. For some aquatic (especially vertebrate) species, there seems to be a certain consistency in the microbiome, e.g. the presence of *Proteobacteria*, *Fusobacteria* and *Firmicutes* in gut microbiomes of zebrafish *Danio rerio* and many other fishes (Roeselers et al. 2011, Ghanbari et al. 2015). For many other aquatic organisms, however, no consistent patterns have thus far been recorded. For *Daphnia*, some bacterial strains (e.g. *Limnohabitans* sp.) have been persistently found in lab cultures (Peerakietkhajorn et al. 2015), but *Daphnia* also associates with oppor-

tunistic species from the natural environment depending on particular conditions (Eckert & Penththaler 2014, Callens et al. 2020). Nevertheless, that does not exclude the possibility that adaptive patterns cannot be found across field and lab microbiomes given the strong genotype × microbiome interactions detected (Macke et al. 2017a, Houwenhuysen et al. 2021). It is likely that there are multiple ‘healthy’ microbial profiles depending on the context. Although the diversity of microbes may be different between field and lab conditions, the functional profile of these communities may be similar (Burke et al. 2011, Adamovsky et al. 2018).

In conclusion, well-designed experiments covering the vast aquatic diversity will greatly improve our mechanistic understanding of which microbial symbionts improve the health of hosts. These insights will allow for the design of powerful models to project the multitude of anthropogenic impacts on aquatic organisms and ecosystems. Validation of such models will require more *in situ*, observational data. Long-term data series that span periods of changing anthropogenic influences on environmental and organism-associated microbiomes will be essential to substantiate and fine-tune model projections, in order to enable the development of adequate measures to counteract the potential negative consequences for organismic and environmental health.

Acknowledgements. W.S. was funded by the FWO Flanders (1252821N). R.S. was supported by BRAIN-be 2.0 under the MicroResist project (B2/191/P1/MicroResist). C.L. has been funded by the Alexander von Humboldt Foundation. C.T. is a PhD fellow of the KU Leuven. V.D. is supported by a post-doctoral mandate (PDM) from KU Leuven. E.D. was funded by KU Leuven research project C16/17/002 and the FWO projects G092619N and G060216N. H.P.G. was supported by the DFG-Dynatriat-project GR1540/30-1.

LITERATURE CITED

- Aalto SL, Decaestecker E, Pulkkinen K (2015) A three-way perspective of stoichiometric changes on host–parasite interactions. *Trends Parasitol* 31:333–340
- Adair KL, Douglas AE (2017) Making a microbiome: the many determinants of host-associated microbial community composition. *Curr Opin Microbiol* 35:23–29
- Adamovsky O, Buerger AN, Wormington AM, Ector N, Griffitt RJ, Bisesi JH Jr, Martyniuk CJ (2018) The gut microbiome and aquatic toxicology: an emerging concept for environmental health. *Environ Toxicol Chem* 37: 2758–2775
- Akbar S, Gu L, Sun Y, Zhou Q and others (2020) Changes in the life history traits of *Daphnia magna* are associated with the gut microbiota composition shaped by diet and antibiotics. *Sci Total Environ* 705:135827
- Almeida AR, Alves M, Domingues I, Henriques I (2019) The

- impact of antibiotic exposure in water and zebrafish gut microbiomes: a 16S rRNA gene-based metagenomic analysis. *Ecotoxicol Environ Saf* 186:109771
- Amalfitano S, Coci M, Corno G, Luna GM (2015) A microbial perspective on biological invasions in aquatic ecosystems. *Hydrobiologia* 746:13–22
- Amin SA, Green DH, Hart MC, Küpper FC, Sunda WG, Carrano CJ (2009) Photolysis of iron–siderophore chelates promotes bacterial–algal mutualism. *Proc Natl Acad Sci USA* 106:17071–17076
- Amin SA, Hmelo LR, Van Tol HM, Durham BP and others (2015) Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* 522: 98–101
- Baker DM, Freeman CJ, Wong JC, Fogel ML, Knowlton N (2018) Climate change promotes parasitism in a coral symbiosis. *ISME J* 12:921–930
- Baquiran JIP, Conaco C (2018) Sponge–microbe partnerships are stable under eutrophication pressure from mariculture. *Mar Pollut Bull* 136:125–134
- Bass D, Stentiford GD, Wang HC, Koskella B, Tyler CR (2019) The pathobiome in animal and plant diseases. *Trends Ecol Evol* 34:996–1008
- Bates JM, Mittge E, Kuhlman J, Baden KN, Cheesman SE, Guillemin K (2006) Distinct signals from the microbiota promote different aspects of zebrafish gut differentiation. *Dev Biol* 297:374–386
- Becker MH, Brucker RM, Schwantes CR, Harris RN, Minbiole KP (2009) The bacterially produced metabolite violacein is associated with survival of amphibians infected with a lethal fungus. *Appl Environ Microbiol* 75:6635–6638
- Berg G, Rybakova D, Fischer D, Cernava T and others (2020) Microbiome definition re-visited: old concepts and new challenges. *Microbiome* 8:103
- Bestion E, Soriano-Redondo A, Cucherousset J, Jacob S and others (2019) Altered trophic interactions in warming climates: consequences for predator diet breadth and fitness. *Proc R Soc B* 286:20192227
- Bijnens K, Thijs S, Leynen N, Stevens V and others (2021) Differential effect of silver nanoparticles on the microbiome of adult and developing planaria. *Aquat Toxicol* 230:105672
- Bilal M, Mehmood S, Rasheed T, Iqbal HMN (2020) Antibiotics traces in the aquatic environment: persistence and adverse environmental impact. *Curr Opin Environ Sci Health* 13:68–74
- Bindoff NL, Cheung WWL, Kairo JG, Arstegui J and others (2019) Changing ocean, marine ecosystems, and dependent communities. In: Pörtner HO, Roberts DC, Masson-Delmotte V, Zhai P and others (eds) IPCC special report on the ocean and cryosphere in a changing climate. IPCC, Geneva, p 447–587
- Björk JR, Diéz-Vives C, Astudillo-García C, Archie EA, Montoya JM (2019) Vertical transmission of sponge microbiota is inconsistent and unfaithful. *Nat Ecol Evol* 3:1172–1183
- Bojarski B, Kot B, Witeska M (2020) Antibacterials in aquatic environment and their toxicity to fish. *Pharmaceuticals (Basel)* 13:189
- Boudry A, Devliegere S, Houwenhuysse S, Clarysse L, Macke E, Vanoverberghe I, Decaestecker E (2020) *Daphnia magna* tolerance to toxic cyanobacteria in the presence of an opportunistic infection within an evolutionary perspective. *Belg J Zool* 150:81–93
- Bourne DG, Morrow KM, Webster NS (2016) Insights into the coral microbiome: underpinning the health and resilience of reef ecosystems. *Annu Rev Microbiol* 70:317–340
- Bright M, Bulgheresi S (2010) A complex journey: transmission of microbial symbionts. *Nat Rev Microbiol* 8:218–230
- Burke C, Steinberg P, Rusch D, Kjelleberg S, Thomas T (2011) Bacterial community assembly based on functional genes rather than species. *Proc Natl Acad Sci USA* 108:14288–14293
- Byrne AQ, Vredenburg VT, Martel A, Pasmans F and others (2019) Cryptic diversity of a widespread global pathogen reveals expanded threats to amphibian conservation. *Proc Natl Acad Sci USA* 116:20382–20387
- Callens M, Macke E, Muylaert K, Bossier P, Lievens B, Waud M, Decaestecker E (2016) Food availability affects the strength of mutualistic host–microbiota interactions in *Daphnia magna*. *ISME J* 10:911–920
- Callens M, Watanabe H, Kato Y, Miura J, Decaestecker E (2018) Microbiota inoculum composition affects holobiont assembly and host growth in *Daphnia*. *Microbiome* 6:56
- Callens M, De Meester L, Muylaert K, Mukherjee S, Decaestecker E (2020) The bacterioplankton community composition and a host genotype dependent occurrence of taxa shape the *Daphnia magna* gut bacterial community. *FEMS Microbiol Ecol* 96:fiaa128
- Carrier TJ, Wolfe K, Lopez K, Gall M, Janies DA, Byrne M, Reitzel AM (2018) Diet-induced shifts in the crown-of-thorns (*Acanthaster* sp.) larval microbiome. *Mar Biol* 165: 157
- Cavicchioli R, Ripple WJ, Timmis KN, Azam F and others (2019) Scientists' warning to humanity: microorganisms and climate change. *Nat Rev Microbiol* 17:569–586
- Chan CX, Vaysberg P, Price DC, Pelletreau KN, Rumpho ME, Bhattacharya D (2018) Active host response to algal symbionts in the sea slug *Elysia chlorotica*. *Mol Biol Evol* 35:1706–1711
- Chaturvedi AD, Pal D, Penta S, Kumar A (2015) Ecotoxic heavy metals transformation by bacteria and fungi in aquatic ecosystem. *World J Microbiol Biotechnol* 31: 1595–1603
- Christie-Oleza JA, Sousoni D, Lloyd M, Armengaud J, Scanlan DJ (2017) Nutrient recycling facilitates long-term stability of marine microbial phototroph–heterotroph interactions. *Nat Microbiol* 2:17100
- Cirimotich CM, Dong Y, Clayton AM, Sandiford SL, Souza-Neto JA, Mulenga M, Dimopoulos G (2011) Natural microbe-mediated refractoriness to *Plasmodium* infection in *Anopheles gambiae*. *Science* 332:855–858
- Cirri E, De Decker S, Bilcke G, Werner M and others (2019) Associated bacteria affect sexual reproduction by altering gene expression and metabolic processes in a biofilm inhabiting diatom. *Front Microbiol* 10:1970
- Clay K (2014) Defensive symbiosis: a microbial perspective. *Funct Ecol* 28:293–298
- Cohen ML, Mashanova EV, Rosen NM, Soto W (2019) Adaptation to temperature stress by *Vibrio fischeri* facilitates this microbe's symbiosis with the Hawaiian bobtail squid (*Euprymna scolopes*). *Evolution* 73:1885–1897
- Cohen ML, Mashanova EV, Jagannathan SV, Soto W (2020) Adaptation to pH stress by *Vibrio fischeri* can affect its symbiosis with the Hawaiian bobtail squid (*Euprymna scolopes*). *Microbiology* 166:262–277
- Coopman M, Muylaert K, Lange B, Reyserhove L, Decaestecker E (2014) Context dependency of infectious disease: the cyanobacterium *Microcystis aeruginosa* decreases white bacterial disease in *Daphnia magna*. *Freshw Biol* 59:714–723

- Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG (2005) Algae acquire vitamin B₁₂ through a symbiotic relationship with bacteria. *Nature* 438:90–93
- Dahan D, Jude BA, Lamendella R, Keesing F, Perron GG (2018) Exposure to arsenic alters the microbiome of larval zebrafish. *Front Microbiol* 9:1323
- Darchambeau F, Faerøvig PJ, Hessen DO (2003) How *Daphnia* copes with excess carbon in its food. *Oecologia* 136: 336–346
- Daskin JH, Alford RA (2012) Context-dependent symbioses and their potential roles in wildlife diseases. *Proc R Soc B* 279:1457–1465
- Davidson AM, Jennions M, Nicotra AB (2011) Do invasive species show higher phenotypic plasticity than native species and, if so, is it adaptive? A meta-analysis. *Ecol Lett* 14: 419–431
- Davoodi S, Foley E (2020) Host-microbe-pathogen interactions: a review of *Vibrio cholerae* pathogenesis in *Drosophila*. *Front Immunol* 10:3128
- De Bary A (1879) Die Erscheinung der Symbiose: Vortrag gehalten auf der Versammlung deutscher Naturforscher und Ärzte zu Cassel. Trübner, Strassburg
- De Meester L, Brans KI, Govaert L, Souffreau C and others (2019) Analysing eco-evolutionary dynamics—the challenging complexity of the real world. *Funct Ecol* 33:43–59
- Decelle J, Colin S, Foster RA (2015) Photosymbiosis in marine planktonic protists. In: Ohtsuka S, Suzuki T, Horiguchi T, Suzuki N, Not F (eds) *Marine protists: diversity and dynamics*. Springer, Tokyo, p 465–500
- Dehler CE, Secombes CJ, Martin SA (2017) Seawater transfer alters the intestinal microbiota profiles of Atlantic salmon (*Salmo salar* L.). *Sci Rep* 7:13877
- Deines P, Bosch TCG (2016) Transitioning from microbiome composition to microbial community interactions: the potential of the metaorganism *Hydra* as an experimental model. *Front Microbiol* 7:1610
- Dheilly NM, Poulin R, Thomas F (2015) Biological warfare: microorganisms as drivers of host–parasite interactions. *Infect Genet Evol* 34:251–259
- Diaz RJ, Rosenberg R (2008) Spreading dead zones and consequences for marine ecosystems. *Science* 321:926–929
- Dirksen P, Assié A, Zimmermann J, Zhang F and others (2020) CeMbio—the *Caenorhabditis elegans* microbiome resource. *G3 (Bethesda)* 10:3025–3039
- Dittami SM, Duboscq-Bidot L, Perennou M, Gobet A, Corre E, Boyen C, Tonon T (2016) Host–microbe interactions as a driver of acclimation to salinity gradients in brown algal cultures. *ISME J* 10:51–63
- Dittami SM, Arboleda E, Auguet JC, Bigalke A and others (2021) A community perspective on the concept of marine holobionts: current status, challenges, and future directions. *PeerJ* 9:e10911
- Dominguez-Bello MG, Godoy-Vitorino F, Knight R, Blaser MJ (2019) Role of the microbiome in human development. *Gut* 68:1108–1114
- Doney SC, Busch DS, Cooley SR, Kroeker KJ (2020) The impacts of ocean acidification on marine ecosystems and reliant human communities. *Annu Rev Environ Resour* 45:83–112
- Dong M, Feng L, Kuang SY, Liu Y and others (2013) Growth, body composition, intestinal enzyme activities and microflora of juvenile Jian carp (*Cyprinus carpio* var. Jian) fed graded levels of dietary valine. *Aquacult Nutr* 19:1–14
- Douglas AE (2018) Omics and the metabolic function of insect–microbial symbioses. *Curr Res Insect Sci* 29:1–6
- Dubilier N, Bergin C, Lott C (2008) Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nat Rev Microbiol* 6:725–740
- Duffy MA, Ochs JH, Penczykowski RM, Civitello DJ, Klausmeier CA, Hall SR (2012) Ecological context influences epidemic size and parasite-driven evolution. *Science* 335:1636–1638
- Dugan HA, Bartlett SL, Burke SM, Doubek JP and others (2017) Salting our freshwater lakes. *Proc Natl Acad Sci USA* 114:4453–4458
- Dunlap PV, Ast JC, Kimura S, Fukui A, Yoshino T, Endo H (2007) Phylogenetic analysis of host–symbiont specificity and codivergence in bioluminescent symbioses. *Cladistics* 23:507–532
- Ebert D (2013) The epidemiology and evolution of symbionts with mixed-mode transmission. *Annu Rev Ecol Evol Syst* 44:623–643
- Eckert EM, Pernthaler J (2014) Bacterial epibionts of *Daphnia*: a potential route for the transfer of dissolved organic carbon in freshwater food webs. *ISME J* 8:1808–1819
- Eckert EM, Anicic N, Fontaneto D (2021) Freshwater zooplankton microbiome composition is highly flexible and strongly influenced by the environment. *Mol Ecol* 30: 1545–1558
- Egerton S, Culloty S, Whooley J, Stanton C, Ross RP (2018) The gut microbiota of marine fish. *Front Microbiol* 9:873
- Englert D, Zubrod JP, Schulz R, Bundschuh M (2013) Effects of municipal wastewater on aquatic ecosystem structure and function in the receiving stream. *Sci Total Environ* 454–455:401–410
- Evariste L, Barret M, Mottier A, Mouchet F, Gauthier L, Pinelli E (2019) Gut microbiota of aquatic organisms: a key endpoint for ecotoxicological studies. *Environ Pollut* 248: 989–999
- Falony G, Joossens M, Vieira-Silva S, Wang J and others (2016) Population-level analysis of gut microbiome variation. *Science* 352:560–564
- Fan L, Liu M, Simister R, Webster NS, Thomas T (2013) Marine microbial symbiosis heats up: the phylogenetic and functional response of a sponge holobiont to thermal stress. *ISME J* 7:991–1002
- Ferrario F, Iveša L, Jaklin A, Perkol-Finkel S, Airoldi L (2016) The overlooked role of biotic factors in controlling the ecological performance of artificial marine habitats. *J Appl Ecol* 53:16–24
- Fieth RA, Gauthier MEA, Bayes J, Green KM, Degnan SM (2016) Ontogenetic changes in the bacterial symbiont community of the tropical demosponge *Amphimedon queenslandica*: metamorphosis is a new beginning. *Front Mar Sci* 3:228
- Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA (2017) The evolution of host–symbiont dependence. *Nat Commun* 8:15973
- Flandroy L, Pouthaidis T, Berg G, Clarke G and others (2018) The impact of human activities and lifestyles on the interlinked microbiota and health of humans and of ecosystems. *Sci Total Environ* 627:1018–1038
- Flórez LV, Biedermann PHW, Engl T, Kaltenpoth M (2015) Defensive symbioses of animals with prokaryotic and eukaryotic microorganisms. *Nat Prod Rep* 32:904–936
- Fontaine SS, Kohl KD (2020) Gut microbiota of invasive bullfrog tadpoles responds more rapidly to temperature than a noninvasive congener. *Mol Ecol* 29:2449–2462
- Fontaine SS, Novarro AJ, Kohl KD (2018) Environmental temperature alters the digestive performance and gut

- microbiota of a terrestrial amphibian. *J Exp Biol* 221: jeb187559
- Foster RA, Zehr JP (2019) Diversity, genomics, and distribution of phytoplankton–cyanobacterium single-cell symbiotic associations. *Annu Rev Microbiol* 73:435–456
- Frankel-Bricker J, Song MJ, Benner MJ, Schaack S (2020) Variation in the microbiota associated with *Daphnia magna* across genotypes, populations, and temperature. *Microb Ecol* 79:731–742
- Freudenthal HD (1962) *Symbiodinium* gen. nov. and *Symbiodinium microadriaticum* sp. nov., a zooxanthella: taxonomy, life cycle, and morphology. *J Protozool* 9:45–52
- Frischkorn KR, Rouco M, Van Mooy BAS, Dyhrman ST (2017) Epibionts dominate metabolic functional potential of *Trichodesmium* colonies from the oligotrophic ocean. *ISME J* 11:2090–2101
- Funkhouser LJ, Bordenstein SR (2013) Mom knows best: the universality of maternal microbial transmission. *PLOS Biol* 11:e1001631
- Gandon S, Capowiez Y, Dubois Y, Michalakis Y, Olivieri I (1996) Local adaptation and gene-for-gene coevolution in a metapopulation model. *Proc R Soc B* 263:1003–1009
- Gegner HM, Ziegler M, Rådecker N, Buitrago-López C, Aranda M, Voolstra CR (2017) High salinity conveys thermotolerance in the coral model *Aiptasia*. *Biol Open* 6:1943–1948
- Ghanbari M, Kneifel W, Domig KJ (2015) A new view of the fish gut microbiome: advances from next-generation sequencing. *Aquaculture* 448:464–475
- Giger W, Alder AC, Golet EM, Kohler HPE and others (2003) Occurrence and fate of antibiotics as trace contaminants in wastewaters, sewage sludges, and surface waters. *Chimia (Aarau)* 57:485–491
- Glynn PW (1991) Coral reef bleaching in the 1980s and possible connections with global warming. *Trends Ecol Evol* 6:175–179
- Gochfeld DJ, Eason CG, Freeman CJ, Thacker RW, Olson JB (2012) Disease and nutrient enrichment as potential stressors on the Caribbean sponge *Aplysina cauliformis* and its bacterial symbionts. *Mar Ecol Prog Ser* 456: 101–111
- Goetsch W (1924) Die Symbiose der Süßwasser-Hydroiden und ihre künstliche Beeinflussung. *Z Morphol Oekol Tiere* 1:660–751
- Goffredi SK, Tilic E, Mullin SW, Dawson KS and others (2020) Methanotrophic bacterial symbionts fuel dense populations of deep-sea feather duster worms (Sabellida, Annelida) and extend the spatial influence of methane seepage. *Sci Adv* 6:eaay8562
- Gordon BR, Leggat W (2010) *Symbiodinium*–invertebrate symbioses and the role of metabolomics. *Mar Drugs* 8: 2546–2568
- Gorokhova E, Rivetti C, Furuhausen S, Edlund A, Ek K, Breitholtz M (2015) Bacteria-mediated effects of antibiotics on *Daphnia* nutrition. *Environ Sci Technol* 49:5779–5787
- Gothwal R, Shashidhar T (2015) Antibiotic pollution in the environment: a review. *Clean (Weinh)* 43:479–489
- Graça AP, Bondoso J, Gaspar H, Xavier JR and others (2013) Antimicrobial activity of heterotrophic bacterial communities from the marine sponge *Erylus discophorus* (Astrophorida, Geodiidae). *PLOS ONE* 8:e78992
- Greenspan SE, Lyra ML, Migliorini GH, Kersch-Becker MF and others (2019) Arthropod–bacteria interactions influence assembly of aquatic host microbiome and pathogen defense. *Proc R Soc B* 286:20190924
- Grosberg RK, Vermeij GJ, Wainwright PC (2012) Biodiversity in water and on land. *Curr Biol* 22:R900–R903
- Grossart HP, Riemann L, Tang KW (2013) Molecular and functional ecology of aquatic microbial symbionts. *Front Microbiol* 4:59
- Haag CR, Ebert D (2004) Parasite-mediated selection in experimental metapopulations of *Daphnia magna*. *Proc R Soc B* 271:2149–2155
- Hardivillier Y, Leignel V, Denis F, Uguen G, Cosson R, Laulier M (2004) Do organisms living around hydrothermal vent sites contain specific metallothioneins? The case of the genus *Bathymodiolus* (Bivalvia, Mytilidae). *Comp Biochem Physiol C Toxicol Pharmacol* 139:111–118
- Harris JM (1993) The presence, nature, and role of gut microflora in aquatic invertebrates: a synthesis. *Microb Ecol* 25:195–231
- Harris RN, Brucker RM, Walke JB, Becker MH and others (2009) Skin microbes on frogs prevent morbidity and mortality caused by a lethal skin fungus. *ISME J* 3:818–824
- Harrison XA, Price SJ, Hopkins K, Leung W, Sergeant C, Garner TW (2019) Diversity–stability dynamics of the amphibian skin microbiome and susceptibility to a lethal viral pathogen. *Front Microbiol* 10:2883
- He S, Wang Q, Li S, Ran C, Guo X, Zhang Z, Zhou Z (2017) Antibiotic growth promoter olaquinox increases pathogen susceptibility in fish by inducing gut microbiota dysbiosis. *Sci China Life Sci* 60:1260–1270
- He J, Lange J, Marinos G, Bathia J and others (2020) Advancing our functional understanding of host–microbiota interactions: a need for new types of studies. *BioEssays* 42:e1900211
- Heard MJ, Smith KF, Ripp KJ, Berger M and others (2013) The threat of disease increases as species move toward extinction. *Conserv Biol* 27:1378–1388
- Hellinger J, Jägers P, Donner M, Sutt F and others (2017) The flashlight fish *Anomalops katoptron* uses bioluminescent light to detect prey in the dark. *PLOS ONE* 12: e0170489
- Herlemann DPR, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF (2011) Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J* 5:1571–1579
- Hinzke T, Kleiner M, Breusing C, Felbeck H and others (2019) Host–microbe interactions in the chemosynthetic *Riftia pachyptila* symbiosis. *MBio* 10:e02243-19
- Holt CC, Bass D, Stentiford GD, van der Giezen M (in press) Understanding the role of the shrimp gut microbiome in health and disease. *J Invertebr Pathol*, <https://doi.org/10.1016/j.jip.2020.107387>
- Hooper LV, Littman DR, Macpherson AJ (2012) Interactions between the microbiota and the immune system. *Science* 336:1268–1273
- Hörtnagl PH, Sommaruga R (2007) Photo-oxidative stress in symbiotic and aposymbiotic strains of the ciliate *Paramecium bursaria*. *Photochem Photobiol Sci* 6:842–847
- Houwenhuyse S, Macke E, Reyserhove L, Bulteel L, Decaestecker E (2018) Back to the future in a petri dish: origin and impact of resurrected microbes in natural populations. *Evol Appl* 11:29–41
- Houwenhuyse S, Stoks R, Mukherjee S, Decaestecker E (2021) Locally adapted gut microbiomes mediate host stress tolerance. *ISME J* 15:2401–2414
- Huntingford C, Williamson MS, Nijse FJMM (2020) CMIP6 climate models imply high committed warming. *Clim Change* 162:1515–1520

- ✦ Huot C, Clerissi C, Gourbal B, Galinier R, Duval D, Toulza E (2020) Schistosomiasis vector snails and their microbiota display a phyllosymbiosis pattern. *Front Microbiol* 10:3092
- ✦ Huyben D, Sun L, Moccia R, Kiessling A, Dicksved J, Lundh T (2018) Dietary live yeast and increased water temperature influence the gut microbiota of rainbow trout. *J Appl Microbiol* 124:1377–1392
- ✦ Jal S, Khora SS (2015) An overview on the origin and production of tetrodotoxin, a potent neurotoxin. *J Appl Microbiol* 119:907–916
- ✦ Jani AJ, Briggs CJ (2018) Host and aquatic environment shape the amphibian skin microbiome but effects on downstream resistance to the pathogen *Batrachochytrium dendrobatidis* are variable. *Front Microbiol* 9:487
- ✦ Jannasch HW (1985) Review lecture: the chemosynthetic support of life and the microbial diversity at deep-sea hydrothermal vents. *Proc R Soc B* 225:277–297
- ✦ Jannasch HW, Mottl MJ (1985) Geomicrobiology of deep-sea hydrothermal vents. *Science* 229:717–725
- ✦ Kan H, Zhao F, Zhang XX, Ren H, Gao S (2015) Correlations of gut microbial community shift with hepatic damage and growth inhibition of *Carassius auratus* induced by pentachlorophenol exposure. *Environ Sci Technol* 49:11894–11902
- ✦ Karntanut W, Pascoe D (2005) Effects of removing symbiotic green algae on the response of *Hydra viridissima* (Pallas 1776) to metals. *Ecotoxicol Environ Saf* 60:301–305
- ✦ Kashyap PC, Marcobal A, Ursell LK, Larauche M and others (2013) Complex interactions among diet, gastrointestinal transit, and gut microbiota in humanized mice. *Gastroenterology* 144:967–977
- ✦ Kayath CA, Ibala Zamba A, Goma-Tchimbakala J, Mamonékéné V, Mombo Makanga GM, Lebonguy AA, Nguimbi E (2019) Microbiota landscape of gut system of guppy fish (*Poecilia reticulata*) plays an outstanding role in adaptation mechanisms. *Int J Microbiol* 2019:3590584
- ✦ Keerthisinghe TP, Wang F, Wang M, Yang Q and others (2020) Long-term exposure to TET increases body weight of juvenile zebrafish as indicated in host metabolism and gut microbiome. *Environ Int* 139:105705
- ✦ Kimura K, Tomaru Y (2014) Coculture with marine bacteria confers resistance to complete viral lysis of diatom cultures. *Aquat Microb Ecol* 73:69–80
- ✦ Kirk D, Jones N, Peacock S, Phillips J, Molnár PK, Krkošek M, Luijckx P (2018) Empirical evidence that metabolic theory describes the temperature dependency of within-host parasite dynamics. *PLOS Biol* 16:e2004608
- ✦ Kivistik C, Knobloch J, Käiro K, Tammert H, Kisand V, Hildebrandt JP, Herlemann DPR (2020) Impact of salinity on the gastrointestinal bacterial community of *Theodoxus fluviatilis*. *Front Microbiol* 11:683
- ✦ Knutie SA, Wilkinson CL, Kohl KD, Rohr JR (2017) Early-life disruption of amphibian microbiota decreases later-life resistance to parasites. *Nat Commun* 8:86
- ✦ Koedooder C, Stock W, Willems A, Mangelinckx S, De Troch M, Vyverman W, Sabbe K (2019) Diatom–bacteria interactions modulate the composition and productivity of benthic diatom biofilms. *Front Microbiol* 10:1255
- ✦ Kohl KD, Yahn J (2016) Effects of environmental temperature on the gut microbial communities of tadpoles. *Environ Microbiol* 18:1561–1565
- ✦ Kohl KD, Weiss RB, Cox J, Dale C, Dearing MD (2014) Gut microbes of mammalian herbivores facilitate intake of plant toxins. *Ecol Lett* 17:1238–1246
- ✦ Koskella B, Bergelson J (2020) The study of host–microbiome (co)evolution across levels of selection. *Philos Trans R Soc B* 375:20190604
- ✦ Koskella B, Hall LJ, Metcalf CJE (2017) The microbiome beyond the horizon of ecological and evolutionary theory. *Nat Ecol Evol* 1:1606–1615
- ✦ Krediet CJ, Ritchie KB, Alagely A, Teplitski M (2013) Members of native coral microbiota inhibit glycosidases and thwart colonization of coral mucus by an opportunistic pathogen. *ISME J* 7:980–990
- ✦ Krotman Y, Yergaliyev TM, Shani RA, Avrahami Y, Szitenberg A (2020) Dissecting the factors shaping fish skin microbiomes in a heterogeneous inland water system. *Microbiome* 8:9
- ✦ Kurth C, Wasmuth I, Wichard T, Pohnert G, Nett M (2019) Algae induce siderophore biosynthesis in the freshwater bacterium *Cupriavidus necator* H16. *Biomaterials* 32:77–88
- ✦ Lachnit T, Bosch TCG, Deines P (2019) Exposure of the host-associated microbiome to nutrient-rich conditions may lead to dysbiosis and disease development—an evolutionary perspective. *MBio* 10:e00355-19
- ✦ Laine AL, Hanski I (2006) Large-scale spatial dynamics of a specialist plant pathogen in a fragmented landscape. *J Ecol* 94:217–226
- ✦ Lan Y, Sun J, Zhang W, Xu T and others (2019) Host–symbiont interactions in deep-sea chemosymbiotic vesicomyid clams: insights from transcriptome sequencing. *Front Mar Sci* 6:680
- ✦ Laspoumaderes C, Modenutti B, Elser JJ, Balseiro E (2015) Does the stoichiometric carbon:phosphorus knife edge apply for predaceous copepods? *Oecologia* 178:557–569
- ✦ Lavy A, Keren R, Yahel G, Ilan M (2016) Intermittent hypoxia and prolonged suboxia measured *in situ* in a marine sponge. *Front Mar Sci* 3:263
- ✦ Lee JJ (2006) Algal symbiosis in larger foraminifera. *Symbiosis* 42:63–75
- ✦ Leger E, Webster JP (2017) Hybridizations within the genus *Schistosoma*: implications for evolution, epidemiology and control. *Parasitology* 144:65–80
- ✦ Leibold MA, Holyoak M, Mouquet N, Amarasekare P and others (2004) The metacommunity concept: a framework for multi-scale community ecology. *Ecol Lett* 7:601–613
- ✦ Leles SG, Mitra A, Flynn KJ, Stoecker DK and others (2017) Oceanic protists with different forms of acquired phototrophy display contrasting biogeographies and abundance. *Proc R Soc B* 284:20170664
- ✦ Levin L, Ekau W, Gooday A, Jorissen F and others (2009) Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences* 6:2063–2098
- ✦ Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI (2008) Worlds within worlds: evolution of the vertebrate gut microbiota. *Nat Rev Microbiol* 6:776–788
- ✦ Li YF, Yang N, Liang X, Yoshida A and others (2018) Elevated seawater temperatures decrease microbial diversity in the gut of *Mytilus coruscus*. *Front Physiol* 9:839
- ✦ Li YF, Xu JK, Chen YW, Ding WY and others (2019) Characterization of gut microbiome in the mussel *Mytilus galloprovincialis* in response to thermal stress. *Front Physiol* 10:1086
- ✦ Limbu SM, Zhou L, Sun SX, Zhang ML, Du ZY (2018) Chronic exposure to low environmental concentrations and legal aquaculture doses of antibiotics cause systemic adverse effects in Nile tilapia and provoke differential human health risk. *Environ Int* 115:205–219
- ✦ Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight

- R (2012) Diversity, stability and resilience of the human gut microbiota. *Nature* 489:220–230
- ✦ Luter HM, Gibb K, Webster NS (2014) Eutrophication has no short-term effect on the *Cymbastela stipitata* holobiont. *Front Microbiol* 5:216
- ✦ Macke E, Callens M, De Meester L, Decaestecker E (2017a) Host–genotype dependent gut microbiota drives zooplankton tolerance to toxic cyanobacteria. *Nat Commun* 8:1608
- ✦ Macke E, Tasiemski A, Massol F, Callens M, Decaestecker E (2017b) Life history and eco-evolutionary dynamics in light of the gut microbiota. *Oikos* 126:508–531
- ✦ Macke E, Callens M, Massol F, Vanoverberghe I, De Meester L, Decaestecker E (2020) Diet and genotype of an aquatic invertebrate affect the composition of free-living microbial communities. *Front Microbiol* 11:380
- ✦ Mallon CA, van Elsas JD, Salles JF (2015) Microbial invasions: the process, patterns, and mechanisms. *Trends Microbiol* 23:719–729
- ✦ Malmqvist B, Rundle S (2002) Threats to the running water ecosystems of the world. *Environ Conserv* 29:134–153
- ✦ Maniero GD, Carey C (1997) Changes in selected aspects of immune function in the leopard frog, *Rana pipiens*, associated with exposure to cold. *J Comp Physiol B* 167:256–263
- ✦ Marton JM, Herbert ER, Craft CB (2012) Effects of salinity on denitrification and greenhouse gas production from laboratory-incubated tidal forest soils. *Wetlands* 32:347–357
- ✦ Marzinelli EM, Zagal CJ, Chapman MG, Underwood AJ (2009) Do modified habitats have direct or indirect effects on epifauna? *Ecology* 90:2948–2955
- Marzinelli EM, Qiu Z, Dafforn KA, Johnston EL, Steinberg PD, Mayer-Pinto M (2018) Coastal urbanisation affects microbial communities on a dominant marine holobiont. *NPJ Biofilms Microbiomes* 4:1
- ✦ Massol F, Macke E, Callens M, Decaestecker E (2021) A methodological framework to analyse determinants of host–microbiota networks, with an application to the relationships between *Daphnia magna*'s gut microbiota and bacterioplankton. *J Anim Ecol* 90:102–119
- ✦ Mayer-Pinto M, Johnston EL, Hutchings PA, Marzinelli EM and others (2015) Sydney Harbour: a review of anthropogenic impacts on the biodiversity and ecosystem function of one of the world's largest natural harbours. *Mar Freshw Res* 66:1088–1105
- ✦ McFall-Ngai MJ (2014) The importance of microbes in animal development: lessons from the squid–vibrio symbiosis. *Annu Rev Microbiol* 68:177–194
- ✦ McFall-Ngai M, Morin JG (1991) Camouflage by disruptive illumination in leiognathids, a family of shallow-water, bioluminescent fishes. *J Exp Biol* 156:119–137
- ✦ Meron D, Atias E, Kruh LI, Elifantz H, Minz D, Fine M, Banin E (2011) The impact of reduced pH on the microbial community of the coral *Acropora eurystoma*. *ISME J* 5:51–60
- ✦ Michiels NK, Seeburger VC, Kalb N, Meadows MG, Anthes N, Mailli AA, Jack CB (2018) Controlled iris radiance in a diurnal fish looking at prey. *R Soc Open Sci* 5:170838
- ✦ Mies M, Sumida PYG, Rådecker N, Voolstra CR (2017) Marine invertebrate larvae associated with *Symbiodinium*: A mutualism from the start? *Front Ecol Evol* 5:56
- ✦ Mihaljevic JR (2012) Linking metacommunity theory and symbiont evolutionary ecology. *Trends Ecol Evol* 27:323–329
- ✦ Milan M, Carraro L, Fariselli P, Martino ME and others (2018) Microbiota and environmental stress: how pollution affects microbial communities in Manila clams. *Aquat Toxicol* 194:195–207
- ✦ Miller CA, Holm HC, Horstmann L, George JC, Fredricks HF, Van Mooy BAS, Apprill A (2020) Coordinated transformation of the gut microbiome and lipidome of bowhead whales provides novel insights into digestion. *ISME J* 14:688–701
- ✦ Moore CJ (2008) Synthetic polymers in the marine environment: a rapidly increasing, long-term threat. *Environ Res* 108:131–139
- ✦ Morel FMM, Price NM (2003) The biogeochemical cycles of trace metals in the oceans. *Science* 300:944–947
- ✦ Morrow KM, Bourne DG, Humphrey C, Botté ES and others (2015) Natural volcanic CO₂ seeps reveal future trajectories for host–microbial associations in corals and sponges. *ISME J* 9:894–908
- ✦ Motone K, Takagi T, Aburaya S, Miura N, Aoki W, Ueda M (2020) A zeaxanthin-producing bacterium isolated from the algal phycosphere protects coral endosymbionts from environmental stress. *MBio* 11:e01019-19
- ✦ Moya A, Ferrer M (2016) Functional redundancy-induced stability of gut microbiota subjected to disturbance. *Trends Microbiol* 24:402–413
- Muller-Parker G, D'elia CF, Cook CB (2015) Interactions between corals and their symbiotic algae. In: Birkeland C (ed) *Coral reefs in the Anthropocene*. Springer, Dordrecht, p 99–116
- ✦ Murdoch CC, Rawls JF (2019) Commensal microbiota regulate vertebrate innate immunity — insights from the zebrafish. *Front Immunol* 10:2100
- Muscatine L (1990) The role of symbiotic algae in carbon and energy flux in reef corals. *Coral Reefs* 25:75–87
- ✦ Neubauer SC (2013) Ecosystem responses of a tidal freshwater marsh experiencing saltwater intrusion and altered hydrology. *Estuaries Coasts* 36:491–507
- ✦ Nicholls RJ, Cazenave A (2010) Sea-level rise and its impact on coastal zones. *Science* 328:1517–1520
- ✦ Nishiguchi MK (2000) Temperature affects species distribution in symbiotic populations of *Vibrio* spp. *Appl Environ Microbiol* 66:3550–3555
- NOAA (2021) Trends in atmospheric carbon dioxide. www.esrl.noaa.gov/gmd/ccgg/trends/weekly.html (accessed 4/5/2021)
- ✦ Noguchi T, Jeon JK, Arakawa O, Sugita H, Deguchi Y, Shida Y, Hashimoto K (1986) Occurrence of tetrodotoxin and anhydrotetrodotoxin in *Vibrio* sp. isolated from the intestines of a xanthid crab, *Atergatis floridus*. *J Biochem* 99:311–314
- ✦ Nougé O, Gallet R, Chevin LM, Lenormand T (2015) Niche limits of symbiotic gut microbiota constrain the salinity tolerance of brine shrimp. *Am Nat* 186:390–403
- ✦ Nowack ECM, Melkonian M (2010) Endosymbiotic associations within protists. *Philos Trans R Soc B* 365:699–712
- ✦ Nyholm SV (2020) In the beginning: egg–microbe interactions and consequences for animal hosts. *Philos Trans R Soc B* 375:20190593
- ✦ Ochsenkühn MA, Röthig T, D'Angelo C, Wiedenmann J, Voolstra CR (2017) The role of floridoside in osmoadaptation of coral-associated algal endosymbionts to high-salinity conditions. *Sci Adv* 3:e1602047
- ✦ Orr JC, Fabry VJ, Aumont O, Bopp L and others (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437:681–686
- ✦ Peerakietkhajorn S, Tsukada K, Kato Y, Matsuura T, Watan-

- abe H (2015) Symbiotic bacteria contribute to increasing the population size of a freshwater crustacean, *Daphnia magna*. *Environ Microbiol Rep* 7:364–372
- ✦ Peng Q, Chang H, Wang R, You Z and others (2019) Potassium sorbate suppresses intestinal microbial activity and triggers immune regulation in zebrafish (*Danio rerio*). *Food Funct* 10:7164–7173
- ✦ Petroni G, Spring S, Schleifer KH, Verni F, Rosati G (2000) Defensive extrusive ectosymbionts of *Euplotidium* (Ciliophora) that contain microtubule-like structures are bacteria related to *Verrucomicrobia*. *Proc Natl Acad Sci USA* 97:1813–1817
- ✦ Phelps D, Brinkman NE, Keely SP, Anneken EM and others (2017) Microbial colonization is required for normal neurobehavioral development in zebrafish. *Sci Rep* 7:11244
- ✦ Phillips ID, Prestie KS (2017) Evidence for substrate influence on artificial substrate invertebrate communities. *Environ Entomol* 46:926–930
- ✦ Portet A, Toulza E, Lokmer A, Huot C, Duval D, Galinier R, Gourbal B (2021) Experimental infection of the *Biomphalaria glabrata* vector snail by *Schistosoma mansoni* parasites drives snail microbiota dysbiosis. *Microorganisms* 9:1084
- ✦ Prazeres M, Renema W (2019) Evolutionary significance of the microbial assemblages of large benthic foraminifera. *Biol Rev Camb Philos Soc* 94:828–848
- ✦ Prazeres M, Ainsworth T, Roberts TE, Pandolfi JM, Leggat W (2017) Symbiosis and microbiome flexibility in calcifying benthic foraminifera of the Great Barrier Reef. *Microbiome* 5:38
- ✦ Provasoli L (1958) Effect of plant hormones on *Ulva*. *Biol Bull (Woods Hole)* 114:375–384
- ✦ Putnam EE, Goodman AL (2020) B vitamin acquisition by gut commensal bacteria. *PLOS Pathog* 16:e1008208
- ✦ Qiao R, Sheng C, Lu Y, Zhang Y, Ren H, Lemos B (2019) Microplastics induce intestinal inflammation, oxidative stress, and disorders of metabolome and microbiome in zebrafish. *Sci Total Environ* 662:246–253
- ✦ Rafaluk-Mohr C, Ashby B, Dahan DA, King KC (2018) Mutual fitness benefits arise during coevolution in a nematode-defensive microbe model. *Evolution Letters* 2:246–256
- ✦ Rawls JF, Mahowald MA, Ley RE, Gordon JI (2006) Reciprocal gut microbiota transplants from zebrafish and mice to germ-free recipients reveal host habitat selection. *Cell* 127:423–433
- Reyne M, Nolan M, McGuiggan H, Aubry A, Emmerson M, Marnell F, Reid N (2021) Artificial agri-environment scheme ponds do not replicate natural environments despite higher aquatic and terrestrial invertebrate richness and abundance. *J Appl Ecol* 58:304–315
- ✦ Reyserhove L, Samaey G, Muylaert K, Coppé V, Van Colen W, Decaestecker E (2017) A historical perspective of nutrient change impact on an infectious disease in *Daphnia*. *Ecology* 98:2784–2798
- ✦ Ridaura VK, Faith JJ, Rey FE, Cheng J and others (2013) Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science* 341:1241214
- ✦ Ridley CP, Bergquist PR, Harper MK, Faulkner DJ, Hooper JNA, Haygood MG (2005) Speciation and biosynthetic variation in four dictyoceratid sponges and their cyanobacterial symbiont, *Oscillatoria spongelliae*. *Chem Biol* 12:397–406
- ✦ Riquelme C, Rojas A, Flores V, Correa JA (1997) Epiphytic bacteria in a copper-enriched environment in northern Chile. *Mar Pollut Bull* 34:816–820
- Robertson SJ, Lemire P, Maughan H, Goethel A and others (2019) Comparison of co-housing and littermate methods for microbiota standardization in mouse models. *Cell Reports* 27:1910–1919
- ✦ Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin K, Rawls JF (2011) Evidence for a core gut microbiota in the zebrafish. *ISME J* 5:1595–1608
- ✦ Ros M, Lacerda MB, Vázquez-Luis M, Masunari S, Guerra-García JM (2016) Studying exotics in their native range: Can introduced fouling amphipods expand beyond artificial habitats? *Biol Invasions* 18:2983–3000
- ✦ Rosati G, Petroni G, Quochi S, Modeo L, Verni F (1999) Epixenosomes: peculiar epibionts of the hypotrich ciliate *Euplotidium itoi* defend their host against predators. *J Eukaryot Microbiol* 46:278–282
- ✦ Röthig T, Ochsenkühn MA, Roik A, van der Merwe R, Voolstra CR (2016) Long-term salinity tolerance is accompanied by major restructuring of the coral bacterial microbiome. *Mol Ecol* 25:1308–1323
- Rousi H, Korpinen S, Bonsdorff E (2019) Brackish-water benthic fauna under fluctuating environmental conditions: the role of eutrophication, hypoxia, and global change. *Front Mar Sci* 6:464
- ✦ Russell SL (2019) Transmission mode is associated with environment type and taxa across bacteria–eukaryote symbioses: a systematic review and meta-analysis. *FEMS Microbiol Lett* 366:fnz013
- ✦ Sabbadin F, Pesante G, Elias L, Besser K and others (2018) Uncovering the molecular mechanisms of lignocellulose digestion in shipworms. *Biotechnol Biofuels* 11:59
- ✦ Sampson TR, Mazmanian SK (2015) Control of brain development, function, and behavior by the microbiome. *Cell Host Microbe* 17:565–576
- ✦ Sánchez-Bayo F, Goka K, Hayasaka D (2016) Contamination of the aquatic environment with neonicotinoids and its implication for ecosystems. *Front Environ Sci* 4:71
- ✦ Schallenberg M, Armstrong A (2004) Assessment of antibiotic activity in surface water of the lower Taieri Plain and impacts on aquatic bacteria in Lake Waipori, South Otago, New Zealand. *NZ J Mar Freshw Res* 38:19–28
- Schellenberg J, Reichert J, Hardt M, Klingelhöfer I and others (2020) The bacterial microbiome of the long-term aquarium cultured high-microbial abundance sponge *Haliclona cnidata*—sustained bioactivity despite community shifts under detrimental conditions. *Front Mar Sci* 7:266
- ✦ Sedorf H, Griffin NW, Ridaura VK, Reyes A and others (2014) Bacteria from diverse habitats colonize and compete in the mouse gut. *Cell* 159:253–266
- ✦ Selvin J, Priya SS, Kiran GS, Thangavelu T, Bai NS (2009) Sponge-associated marine bacteria as indicators of heavy metal pollution. *Microbiol Res* 164:352–363
- ✦ Sengupta S, Chattopadhyay M, Grossart HP (2013) The multifaceted roles of antibiotics and antibiotic resistance in nature. *Front Microbiol* 4:47
- ✦ Seyedsayamdost MR, Case RJ, Kolter R, Clardy J (2011) The Jekyll-and-Hyde chemistry of *Phaeobacter gallaeciensis*. *Nat Chem* 3:331–335
- ✦ Seymour JR, Amin SA, Raina JB, Stocker R (2017) Zooming in on the phycosphere: the ecological interface for phytoplankton–bacteria relationships. *Nat Microbiol* 2:17065
- ✦ Shoemaker KM, McCliment EA, Moisaner PH (2020) Copepod-associated gammaproteobacterial alkaline phosphatases in the North Atlantic Subtropical Gyre. *Front Microbiol* 11:1033

- Silvestre F, Gillardin V, Dorts J (2012) Proteomics to assess the role of phenotypic plasticity in aquatic organisms exposed to pollution and global warming. *Integr Comp Biol* 52:681–694
- Simister R, Taylor MW, Tsai P, Webster N (2012) Sponge-microbe associations survive high nutrients and temperatures. *PLOS ONE* 7:e52220
- Sison-Mangus MP, Mushegian AA, Ebert D (2015) Water fleas require microbiota for survival, growth and reproduction. *ISME J* 9:59–67
- Smith D, Muscatine L, Lewis D (1969) Carbohydrate movement from autotrophs to heterotrophs in parasitic and mutualistic symbiosis. *Biol Rev Camb Philos Soc* 44:17–85
- Sneed JM, Sharp KH, Ritchie KB, Paul VJ (2014) The chemical cue tetrabromopyrrole from a biofilm bacterium induces settlement of multiple Caribbean corals. *Proc R Soc B* 281:20133086
- Sokolovskaya OM, Shelton AN, Taga ME (2020) Sharing vitamins: cobamides unveil microbial interactions. *Science* 369:eaba0165
- Sommaruga R, Whitehead K, Shick JM, Lobban CS (2006) Mycosporine-like amino acids in the zooxanthella-ciliate symbiosis *Maristentor dinoferus*. *Protist* 157:185–191
- Stock W, Blommaert L, De Troch M, Mangelinckx S, Willems A, Vyverman W, Sabbe K (2019a) Host specificity in diatom-bacteria interactions alleviates antagonistic effects. *FEMS Microbiol Ecol* 95:fiz171
- Stock W, Vanelslander B, Rüdiger F, Sabbe K, Vyverman W, Karsten U (2019b) Thermal niche differentiation in the benthic diatom *Cylindrotheca closterium* (Bacillariophyceae) complex. *Front Microbiol* 10:1395
- Stoecker DK, Johnson MD, de Vargas C, Not F (2009) Acquired phototrophy in aquatic protists. *Aquat Microb Ecol* 57:279–310
- Stratil SB, Neulinger SC, Knecht H, Friedrichs AK, Wahl M (2013) Temperature-driven shifts in the epibiotic bacterial community composition of the brown macroalga *Fucus vesiculosus*. *MicrobiologyOpen* 2:338–349
- Sullam KE, Pichon S, Schaer TMM, Ebert D (2018) The combined effect of temperature and host clonal line on the microbiota of a planktonic crustacean. *Microb Ecol* 76:506–517
- Székely AJ, Langenheder S (2014) The importance of species sorting differs between habitat generalists and specialists in bacterial communities. *FEMS Microbiol Ecol* 87:102–112
- Szymańska U, Wiergowski M, Sołtyszewski I, Kuzemko J, Wiergowska G, Woźniak MK (2019) Presence of antibiotics in the aquatic environment in Europe and their analytical monitoring: recent trends and perspectives. *Microchem J* 147:729–740
- Tadesse S (2018) Why are marine ecosystems biologically more diversified than their equivalent terrestrial ecosystems. *Int J Avian & Wildlife Biol* 3:304–305
- Tasiemski A, Massol F, Cuvillier-Hot V, Boidin-Wichlacz C and others (2015) Reciprocal immune benefit based on complementary production of antibiotics by the leech *Hirudo verbana* and its gut symbiont *Aeromonas veronii*. *Sci Rep* 5:17498
- terHorst CP, Zee PC (2016) Eco-evolutionary dynamics in plant-soil feedbacks. *Funct Ecol* 30:1062–1072
- Thompson RCA, Lymbery AJ, Smith A (2010) Parasites, emerging disease and wildlife conservation. *Int J Parasitol* 40:1163–1170
- Tian RM, Zhang W, Cai L, Wong YH, Ding W, Qian PY (2017) Genome reduction and microbe-host interactions drive adaptation of a sulfur-oxidizing bacterium associated with a cold seep sponge. *mSystems* 2:e00184-16
- Tianero MD, Balaich JN, Donia MS (2019) Localized production of defence chemicals by intracellular symbionts of *Haliclona* sponges. *Nat Microbiol* 4:1149–1159
- Timmis K, Cavicchioli R, Garcia JL, Nogales B and others (2019) The urgent need for microbiology literacy in society. *Environ Microbiol* 21:1513–1528
- Toledo R, Fried B (eds) (2011) *Biomphalaria* snails and larval trematodes. Springer-Verlag, New York, NY
- Tran C, Hadfield MG (2011) Larvae of *Pocillopora damicornis* (Anthozoa) settle and metamorphose in response to surface-biofilm bacteria. *Mar Ecol Prog Ser* 433:85–96
- Trevelline BK, Fontaine SS, Hartup BK, Kohl KD (2019) Conservation biology needs a microbial renaissance: a call for the consideration of host-associated microbiota in wildlife management practices. *Proc R Soc B* 286:20182448
- Vaelli PM, Theis KR, Williams JE, O’Connell LA, Foster JA, Eisthen HL (2020) The skin microbiome facilitates adaptive tetrodotoxin production in poisonous newts. *eLife* 9:e53898
- van der Heide T, Govers LL, de Fouw J, Olf H and others (2012) A three-stage symbiosis forms the foundation of seagrass ecosystems. *Science* 336:1432–1434
- van Marken Lichtenbelt WD (1992) Digestion in an ectothermic herbivore, the green iguana (*Iguana iguana*): effect of food composition and body temperature. *Physiol Zool* 65:649–673
- van Oppen MJH, Blackall LL (2019) Coral microbiome dynamics, functions and design in a changing world. *Nat Rev Microbiol* 17:557–567
- Vanhove MPM, Thys S, Decaestecker E, Antoine-Moussiaux N and others (2020) Global change increases zoonotic risk, COVID-19 changes risk perceptions: a plea for urban nature connectedness. *Cities & Health* (in press), <https://doi.org/10.1080/23748834.2020.1805282>
- Venkatesh B, Lu SQ, Dandona N, See SL, Brenner S, Soong TW (2005) Genetic basis of tetrodotoxin resistance in pufferfishes. *Curr Biol* 15:2069–2072
- Verma SC, Miyashiro T (2013) Quorum sensing in the squid-*Vibrio* symbiosis. *Int J Mol Sci* 14:16386–16401
- Villareal TA (1992) Marine nitrogen-fixing diatom-cyanobacteria symbioses. In: Carpenter EJ, Capone DG, Rueter JG (eds) *Marine pelagic cyanobacteria: Trichodesmium and other diazotrophs*. Springer, Dordrecht, p 163–175
- Voss JD, Richardson LL (2006) Nutrient enrichment enhances black band disease progression in corals. *Coral Reefs* 25:569–576
- Wan Z, Wang C, Zhou J, Shen M, Wang X, Fu Z, Jin Y (2019) Effects of polystyrene microplastics on the composition of the microbiome and metabolism in larval zebrafish. *Chemosphere* 217:646–658
- Wang W, Gao H, Jin S, Li R, Na G (2019) The ecotoxicological effects of microplastics on aquatic food web, from primary producer to human: a review. *Ecotoxicol Environ Saf* 173:110–117
- Webster NS, Cobb RE, Negri AP (2008) Temperature thresholds for bacterial symbiosis with a sponge. *ISME J* 2:830–842
- Webster NS, Soo R, Cobb R, Negri AP (2011) Elevated seawater temperature causes a microbial shift on crustose coralline algae with implications for the recruitment of coral larvae. *ISME J* 5:759–770

- Webster NS, Negri A, Botté E, Laffy P and others (2016) Host-associated coral reef microbes respond to the cumulative pressures of ocean warming and ocean acidification. *Sci Rep* 6:19324
- Webster JP, Gower CM, Knowles SCL, Molyneux DH, Fenton A (2016) One health—an ecological and evolutionary framework for tackling neglected zoonotic diseases. *Evol Appl* 9:313–333
- Welden NAC, Cowie PR (2016) Long-term microplastic retention causes reduced body condition in the langoustine, *Nephrops norvegicus*. *Environ Pollut* 218:895–900
- West HH (1981) Chloroplast symbiosis and development of the ascoglossan opisthobranch *Elysia chlorotica*. PhD dissertation, Northeastern University, Boston, MA
- White JF Jr, Torres MS (2009) Defensive mutualism in microbial symbiosis. CRC Press, Boca Raton, FL
- White RA, Callister SJ, Moore RJ, Baker ES, Jansson JK (2016) The past, present and future of microbiome analyses. *Nat Protoc* 11:2049–2053
- Wichard T, Charrier B, Mineur F, Bothwell JH, Clerck OD, Coates JC (2015) The green seaweed *Ulva*: a model system to study morphogenesis. *Front Plant Sci* 6:72
- Wollenberger L, Halling-Sørensen B, Kusk KO (2000) Acute and chronic toxicity of veterinary antibiotics to *Daphnia magna*. *Chemosphere* 40:723–730
- Xia J, Lu L, Jin C, Wang S and others (2018) Effects of short term lead exposure on gut microbiota and hepatic metabolism in adult zebrafish. *Comp Biochem Physiol C Toxicol Pharmacol* 209:1–8
- Xie NB, Feng L, Liu Y, Jiang J and others (2011) Growth, body composition, intestinal enzyme activities and microflora of juvenile Jian carp (*Cyprinus carpio* var. Jian) fed graded levels of dietary phosphorus. *Aquacult Nutr* 17: 645–656
- Ye S, Bhattacharjee M, Siemann E (2020) Stress tolerance alteration in the freshwater cnidarian green hydra (*Hydra viridissima*) via symbiotic algae mutagenesis. *Symbiosis* 82:189–199
- Yim G, Huimi Wang H, Davies Frs J (2007) Antibiotics as signalling molecules. *Philos Trans R Soc B* 362:1195–1200
- Yu K, Li X, Qiu Y, Zeng X and others (2020) Low-dose effects on thyroid disruption in zebrafish by long-term exposure to oxytetracycline. *Aquat Toxicol* 227:105608
- Zalewski A, Wagner ND, Frost PC (2011) Antibiotics affect the growth responses of *Daphnia magna* to poor food quality. *Aquat Ecol* 45:493–504
- Zan J, Li Z, Tianero MD, Davis J, Hill RT, Donia MS (2019) A microbial factory for defensive kahalalides in a tripartite marine symbiosis. *Science* 364:eaaw6732
- Zhou L, Limbu SM, Shen M, Zhai W and others (2018) Environmental concentrations of antibiotics impair zebrafish gut health. *Environ Pollut* 235:245–254
- Zoccarato L, Grossart HP (2019) Relationship between lifestyle and structure of bacterial communities and their functionality in aquatic systems. In: Hurst CJ (ed) The structure and function of aquatic microbial communities. *Advances in environmental microbiology, Vol 7*. Springer International Publishing, Cham, p 13–52

*Editorial responsibility: Paul del Giorgio,
Montreal, Quebec, Canada
Reviewed by: 3 anonymous referees*

Submitted: December 2, 2020

Accepted: May 17, 2021

Proofs received from author(s): August 23, 2021